arXiv:1302.5187v2 [math.RT] 18 Mar 2013

HEARTS OF TWIN COTORSION PAIRS ON EXACT CATEGORIES

YU LIU

Abstract. In the papers of Nakaoka, he introduced the notion of hearts of (twin) cotorsion pair on triangulated categories and showed that they have structures of (semi-) abelian categories. We study in this article a twin cotorsion pair $(\mathcal{S},\mathcal{T}),(\mathcal{U},\mathcal{V})$ on an exact category $\mathcal{B}$ with enough projectives and injectives and introduce a notion of the heart. First we show that its heart is preabelian. Moreover we show the heart of a single cotorsion pair is abelian. These results are analog of Nakaoka's results in triangulated categories. We also consider special cases where the heart has nicer structure. By our results, the heart of a special twin cotorsion pair $(\mathcal{S},\mathcal{T}),(\mathcal{T},\mathcal{V})$, is integral and almost abelian. Finally we show that the Gabriel-Zisman localisation of the heart at the class of regular morphisms is abelian, and moreover it is equivalent to the category of finitely presented modules over a stable subcategory of $\mathcal{B}$.

Contents

1. Introduction 1
2. Preliminaries 3
3. $\underline{\mathcal{H}}$ is preabelian 8
4. Abelianess of the hearts of cotorsion pairs 12
5. $\underline{\mathcal{H}}$ is semi-abelian 13
6. The case where $\underline{\mathcal{H}}$ becomes integral 16
7. The case where $\underline{\mathcal{H}}$ becomes almost abelian 20
8. Existence of enough projectives/injectives 22
9. Localisation on the heart of a special twin cotorsion pair 25
10. Examples 28
Acknowledgments 31
References 31

1. Introduction

The cotorsion pairs were first introduced by Salce in [16], and it has been deeply studied in the representation theory during these years, especially in tilting theory and Cohen-Macaulay modules [1] (see [7] for more recent examples). Recently, the cotorsion pair are also studied in triangulated categories [9], in particular, Nakaoka introduced the notion of hearts of cotorsion pairs and showed that the hearts are abelian category [13]. This is a generalization of the hearts of t-structure in triangulated categories [3] and the quotient of triangulated categories by cluster tilting subcategories [12]. Moreover, he generalized these results to a more general setting called twin cotorsion pairs [14].

The aim of this paper is to extend Nakaoka's results to cotorsion pair in Quillen's exact categories, which plays an important role in representation theory [10]. We consider a *cotorsion pair* in an exact category (see for example [11, A.1]), which is a pair $(\mathcal{U},\mathcal{V})$ of subcategories of an exact category $\mathcal{B}$ satisfying $\operatorname{Ext}^1_{\mathcal{B}}(\mathcal{U},\mathcal{V})=0$ (*i.e.* $\operatorname{Ext}^1_{\mathcal{B}}(U,V)=0$, $\forall U\in\mathcal{U}$ and $\forall V\in\mathcal{V}$) and any $B\in\mathcal{B}$ admits two short exact sequences $V_B\rightarrowtail U_B\twoheadrightarrow B$ and $B\rightarrowtail V^B\twoheadrightarrow U^B$ where $V_B,V^B\in\mathcal{V}$ and $U_B,U^B\in\mathcal{U}$ (see Definition

---

Key words and phrases. exact category, preabelian category, abelian category, cotorsion pair, twin cotorsion pair, heart.

2.3 for more details). Let

$$\mathcal{B}^+ := \{B \in \mathcal{B} \mid U_B \in \mathcal{V}\}, \quad \mathcal{B}^- := \{B \in \mathcal{B} \mid V^B \in \mathcal{U}\}.$$

We define the *heart* of $(\mathcal{U}, \mathcal{V})$ as the quotient category (see Definition 2.8 for more details)

$$\underline{\mathcal{H}} := (\mathcal{B}^+ \cap \mathcal{B}^-)/(\mathcal{U} \cap \mathcal{V}).$$

An important class of exact categories is given by Frobenius categories, which gives most of important triangulated categories appearing in representation theory. Now we state our first main result, which generalizes [13, Theorem 6.4]. We will prove it in Section 4.

Theorem. *Let $(\mathcal{U}, \mathcal{V})$ be a cotorsion pair on an exact category $\mathcal{B}$. Then $\underline{\mathcal{H}}$ is abelian.*

Moreover, following Nakaoka, we consider pairs of cotorsion pairs $(\mathcal{S}, \mathcal{T})$ and $(\mathcal{U}, \mathcal{V})$ in $\mathcal{B}$ such that $\mathcal{S} \subseteq \mathcal{U}$, we also call such a pair a *twin cotorsion pair* (see Definition 2.5 for more details). The notion of hearts is generalized to such pairs (see Definition 2.7 for more details), and our second main result is the following, which generalize [14, Theorem 5.4]. We will first prove $\underline{\mathcal{H}}$ is preabelian in Section 3 and then show the following theorem in Section 5.

Theorem. *Let $(\mathcal{S}, \mathcal{T}), (\mathcal{U}, \mathcal{V})$ be a twin cotorsion pair on $\mathcal{B}$. Then $\underline{\mathcal{H}}$ is semi-abelian.*

The notion of semi-abelian category was introduced by Rump [15], as a special class of preabelian categories (see Definition 5.1). A especially nice class of semi-abelian categories is called integral (see Definition 6.1, and see [15, §2] for examples). Our third main theorem gives sufficient conditions for hearts to be integral. We will show it in Section 6.

Theorem. *Let $(\mathcal{S}, \mathcal{T}), (\mathcal{U}, \mathcal{V})$ be a twin cotorsion pair on $\mathcal{B}$ satisfying*

$$\mathcal{U} \subseteq \mathcal{S} * \mathcal{T},\ \mathcal{P} \subseteq \mathcal{W} \quad \text{or} \quad \mathcal{T} \subseteq \mathcal{U} * \mathcal{V},\ \mathcal{I} \subseteq \mathcal{W}.$$

Then $\underline{\mathcal{H}}$ is integral.

Another nice class of semi-abelian categories is almost abelian categories. For example, any torsion class associated with a tilting module is almost abelian [5]. Our fourth main theorem gives sufficient conditions for hearts to be almost abelian. We will show it in Section 7.

Theorem. *Let $(\mathcal{S}, \mathcal{T}), (\mathcal{U}, \mathcal{V})$ be a twin cotorsion pair on $\mathcal{B}$ satisfying*

$$\mathcal{U} \subseteq \mathcal{T} \text{ or } \mathcal{T} \subseteq \mathcal{U}.$$

Then $\underline{\mathcal{H}}$ is integral and almost abelian.

Finally, we consider a special twin cotorsion pair $(\mathcal{S}, \mathcal{T}), (\mathcal{T}, \mathcal{V})$, note that this is an analog of TTF theory and recollement. Then we have the following theorem which gives a more explicit description of the heart and can be regarded as an analog of [4, Theorem 5.7]. We will prove it in Section 9.

Theorem. *Let $(\mathcal{S}, \mathcal{T}), (\mathcal{U}, \mathcal{V})$ be a twin cotorsion pari on $\mathcal{B}$ such that $\mathcal{T} = \mathcal{U}$. Let R denote the class of regular morphisms in $\mathcal{B}/\mathcal{T}$ and $(\mathcal{B}/\mathcal{T})_R$ denote the localisation of $\underline{\mathcal{H}} = \mathcal{B}/\mathcal{T}$ at R, then*

$$(\mathcal{B}/\mathcal{T})_R \simeq \operatorname{mod}(\Omega\mathcal{S}/\mathcal{P})$$

where $\Omega\mathcal{S}$ consists of objects ΩS such that there exists a short exact sequence

$$\Omega S \rightarrowtail P \twoheadrightarrow S \ (P \in \mathcal{P}, S \in \mathcal{S}).$$

In Section 2, we collect basic material on twin cotorsion pairs on $\mathcal{B}$. In Section 3-7 and 9, we prove our main results. In Section 8, we consider the cases when the heart of a twin cotorsion pair has enough projectives/injectives. In the last section we study some examples of twin cotorsion pairs.

2. Preliminaries

First we briefly review the important properties of exact categories. For more details, we refer to [2]. Let $\mathcal{A}$ be an additive category, we call a pair of morphisms (i, d) a *weak short exact sequence* if i is the kernel of d and d is the cokernel of i. Let $\mathcal{E}$ be a class of weak short exact sequences of $\mathcal{A}$, stable under isomorphisms, direct sums and direct summands. If a weak short exact sequence (i, d) is in $\mathcal{E}$, we call it a *short exact sequence* and denote it by

$$\begin{tikzcd} X \arrow[r, tail, "i"] & Y \arrow[r, two heads, "d"] & Z. \end{tikzcd}$$

We call i an *inflation* and d a *deflation*. The pair $(\mathcal{A}, \mathcal{E})$ (or simply $\mathcal{A}$) is said to be an *exact category* if it satisfies the following properties:

(a) Identity morphisms are inflations and deflations.
(b) The composition of two inflations (resp. deflations) is an inflation (resp. deflation).
(c) If $X \overset{i}{\rightarrowtail} Y \overset{d}{\twoheadrightarrow} Z$ is a short exact sequence, for any morphisms $f : Z' \to Z$ and $g : X \to X'$, there are commutative diagrams

$$\begin{tikzcd} Y' \arrow[r, two heads, "d'"] \arrow[d, "f'"'] \arrow[dr, phantom, "PB"] & Z' \arrow[d, "f"] & X \arrow[r, tail, "i"] \arrow[d, "g"'] \arrow[dr, phantom, "PO"] & Y \arrow[d, "g'"] \\ Y \arrow[r, two heads, "d"'] & Z & X' \arrow[r, tail, "i'"'] & Y' \end{tikzcd}$$

where d' is a deflation and i' is an inflation, the left square being a pull-back and the right being a push-out.

We introduce the following properties of exact category, the proofs of which can be find in [2, §2]:

Proposition 2.1. *Consider a commutative square*

$$\begin{tikzcd} A \arrow[r, tail, "i"] \arrow[d, "f"'] & B \arrow[d, "f'"] \\ A' \arrow[r, tail, "i'"'] & B' \end{tikzcd}$$

in which i and i' are inflations. The following conditions are equivalent:

(a) *The square is a push-out.*
(b) *The sequence* $A \overset{\binom{i}{-f}}{\rightarrowtail} B \oplus A' \overset{(f'\ i')}{\twoheadrightarrow} B'$ *is short exact.*
(c) *The square is both a push-out and a pull-back.*
(d) *The square is a part of a commutative diagram*

$$\begin{tikzcd} A \arrow[r, tail, "i"] \arrow[d, "f"'] & B \arrow[r, two heads] \arrow[d, "f'"] & C \arrow[d, equal] \\ A' \arrow[r, tail, "i'"'] & B' \arrow[r, two heads] & C \end{tikzcd}$$

with short exact rows.

Proposition 2.2. (a) *If* $X \overset{i}{\rightarrowtail} Y \overset{d}{\twoheadrightarrow} Z$ *and* $N \overset{g}{\rightarrowtail} M \overset{f}{\twoheadrightarrow} Y$ *are two short exact sequences, then there is a commutative diagram of short exact sequences*

$$\begin{tikzcd}
N \arrow[r, equal] \arrow[d, tail] & N \arrow[d, tail, "g"] & \\
Q \arrow[r, tail] \arrow[d, two heads] & M \arrow[r, two heads] \arrow[d, two heads, "f"] & Z \arrow[d, equal] \\
X \arrow[r, tail, "i"'] & Y \arrow[r, two heads, "d"'] & Z
\end{tikzcd}$$

where the lower-left square is both a push-out and a pull-back.

(b) *If* $X \overset{i}{\rightarrowtail} Y \overset{d}{\twoheadrightarrow} Z$ *and* $Y \overset{g}{\rightarrowtail} K \overset{f}{\twoheadrightarrow} L$ *are two short exact sequences, then there is a commutative diagram of short exact sequences*

$$\begin{tikzcd}
X \arrow[r, tail, "i"] \arrow[d, equal] & Y \arrow[r, two heads, "d"] \arrow[d, tail, "g"] & Z \arrow[d, tail] \\
X \arrow[r, tail] & K \arrow[r, two heads] \arrow[d, two heads, "f"] & R \arrow[d, two heads] \\
 & L \arrow[r, equal] & L
\end{tikzcd}$$

where the upper-right square is both a push-out and a pull-back.

Let $\mathcal{A}$ be an exact category, an object P is called projective in $\mathcal{A}$ if for any deflation $f : X \to Y$ and any morphism $g : P \to Y$, there exists a morphism $h : P \to X$ such that $g = fh$. $\mathcal{A}$ is said to have enough projectives if for any object $X \in \mathcal{A}$, there is an object P which is projective in $\mathcal{A}$ and a deflation $p : P \to X$. Injective objects and having enough injectives are defined dually.

Throughout this paper, let $\mathcal{B}$ be an exact category with enough projectives and injectives. Let $\mathcal{P}$ (resp. $\mathcal{I}$) be the full subcategory of projectives (resp. injectives) of $\mathcal{B}$. Throughout this article we assume that $\mathcal{B}$ is Krull-Schmidt.

Definition 2.3. Let $\mathcal{U}$ and $\mathcal{V}$ be full additive subcategories of $\mathcal{B}$ which are closed under direct summands. We call $(\mathcal{U}, \mathcal{V})$ a *cotorsion pair* if it satisfies the following conditions:

(a) $\operatorname{Ext}^1_{\mathcal{B}}(\mathcal{U}, \mathcal{V}) = 0$.
(b) For any object $B \in \mathcal{B}$, there exits two short exact sequences

$$V_B \rightarrowtail U_B \twoheadrightarrow B, \quad B \rightarrowtail V^B \twoheadrightarrow U^B$$

satisfying $U_B, U^B \in \mathcal{U}$ and $V_B, V^B \in \mathcal{V}$.

By definition of a cotorsion pair, we can immediately conclude:

Lemma 2.4. *Let* $(\mathcal{U}, \mathcal{V})$ *be a cotorsion pair of* $\mathcal{B}$*, then*

(a) B *belongs to* $\mathcal{U}$ *if and only if* $\operatorname{Ext}^1_{\mathcal{B}}(B, \mathcal{V}) = 0$.
(b) B *belongs to* $\mathcal{V}$ *if and only if* $\operatorname{Ext}^1_{\mathcal{B}}(\mathcal{U}, B) = 0$.
(c) $\mathcal{U}$ *and* $\mathcal{V}$ *are closed under extension.*
(d) $\mathcal{P} \subseteq \mathcal{U}$ *and* $\mathcal{I} \subseteq \mathcal{V}$.

Definition 2.5. A pair of cotorsion pairs $(\mathcal{S}, \mathcal{T})$, $(\mathcal{U}, \mathcal{V})$ on $\mathcal{B}$ is called a *twin cotorsion pair* if it satisfies:

$$\mathcal{S} \subseteq \mathcal{U}.$$

By definition and Lemma 2.4 this condition is equivalent to $\operatorname{Ext}^1_{\mathcal{B}}(\mathcal{S}, \mathcal{V}) = 0$, and also to $\mathcal{V} \subseteq \mathcal{T}$.

Remark 2.6. We also regard a cotorsion pair $(\mathcal{U}, \mathcal{V})$ as a degenerated case of a twin cotorsion pair $(\mathcal{U}, \mathcal{V}), (\mathcal{U}, \mathcal{V})$.

Definition 2.7. For any twin otorsion pairs $(\mathcal{S},\mathcal{T})$, $(\mathcal{U},\mathcal{V})$, put

$$\mathcal{W} := \mathcal{T} \cap \mathcal{U}.$$

(a) $\mathcal{B}^+$ is defined to be the full subcategory of $\mathcal{B}$, consisting of objects B which admits a short exact sequence

$$V_B \rightarrowtail U_B \twoheadrightarrow B$$

where $U_B \in \mathcal{W}$ and $V_B \in \mathcal{V}$.

(b) $\mathcal{B}^-$ is defined to be the full subcategory of $\mathcal{B}$, consisting of objects B which admits a short exact sequence

$$B \rightarrowtail T^B \twoheadrightarrow S^B$$

where $T^B \in \mathcal{W}$ and $S^B \in \mathcal{S}$.

By this definition we get $\mathcal{S} \subseteq \mathcal{U} \subseteq \mathcal{B}^-$ and $\mathcal{V} \subseteq \mathcal{T} \subseteq \mathcal{B}^+$.

Definition 2.8. Let $(\mathcal{S},\mathcal{T})$, $(\mathcal{U},\mathcal{V})$ be a twin cotorsion pair of $\mathcal{B}$, we denote the quotient of $\mathcal{B}$ by $\mathcal{W}$ as $\underline{\mathcal{B}} := \mathcal{B}/\mathcal{W}$. For any morphism $f \in \operatorname{Hom}_{\mathcal{B}}(X,Y)$, we denote its image in $\operatorname{Hom}_{\underline{\mathcal{B}}}(X,Y)$ by $\underline{f}$. And for any subcategory $\mathcal{C}$ of $\mathcal{B}$, we denote by $\underline{\mathcal{C}}$ the subcategory of $\underline{\mathcal{B}}$ consisting of the same objects as $\mathcal{C}$. Put

$$\mathcal{H} := \mathcal{B}^+ \cap \mathcal{B}^-.$$

Since $\mathcal{H} \supseteq \mathcal{W}$, we have an additive full quotient subcategory

$$\underline{\mathcal{H}} := \mathcal{H}/\mathcal{W}$$

which we call the *heart* of twin cotorsion pair $(\mathcal{S},\mathcal{T})$, $(\mathcal{U},\mathcal{V})$.
The heart of a cotorsion pair $(\mathcal{U},\mathcal{V})$ is defined to be the heart of twin cotorsion pair $(\mathcal{U},\mathcal{V})$, $(\mathcal{U},\mathcal{V})$.

We prove some useful lemmas for a twin cotorsion pair $(\mathcal{S},\mathcal{T}),(\mathcal{U},\mathcal{V})$ in the following:

Lemma 2.9. *Let $(\mathcal{S},\mathcal{T}),(\mathcal{U},\mathcal{V})$ be a twin cotorsion pair on $\mathcal{B}$, then*

(a) *$\mathcal{B}^-$ is closed under direct summands. Moreover, if $X \in \mathcal{B}^-$ admits a short exact sequence*

$$X \rightarrowtail W \twoheadrightarrow U$$

where $W \in \mathcal{W}$ and $U \in \mathcal{U}$, then any direct summand X_1 of B admits a short exact sequence

$$X_1 \rightarrowtail W \twoheadrightarrow Y$$

where $Y \in \mathcal{U}$.

(b) *$\mathcal{B}^+$ is closed under direct summands. Moreover, if $X \in \mathcal{B}^+$ admits a short exact sequence*

$$V \rightarrowtail W' \twoheadrightarrow B$$

where $W \in \mathcal{W}$ and $V \in \mathcal{V}$, then any direct summand X_2 of B admits a short exact sequence

$$Z \rightarrowtail W' \twoheadrightarrow X_2$$

where $Z \in \mathcal{V}$.

Proof. We only show (a), (b) is by dual.
Suppose $X_1 \oplus X_2$ admits a short exact sequence

$$\begin{tikzcd} X_1 \oplus X_2 \arrow[r, tail, "(x_1\ x_2)"] & W \arrow[r, two heads] & U \end{tikzcd}$$

where $U \in \mathcal{U}$ and $W \in \mathcal{W}$. Then $x_1 : X_1 \to W$ is also an inflation by the properties of exact category. Let x_1 admits a short exact sequence

$$\begin{tikzcd} X_1 \arrow[r, tail, "x_1"] & W \arrow[r, two heads] & Y. \end{tikzcd}$$

For any morphism $f : X_1 \to V_0$ where $V_0 \in \mathcal{V}$, consider a morphism $\left(\begin{smallmatrix} f & 0 \end{smallmatrix}\right) : X_1 \oplus X_2 \to V_0$. Since $\operatorname{Ext}^1_{\mathcal{B}}(U, V_0) = 0$, $\left(\begin{smallmatrix} x_1 & x_2 \end{smallmatrix}\right)$ is a left $\mathcal{V}$-approximation of W, there exists a morphism $g : W \to V_0$ such that $\left(\begin{smallmatrix} f & 0 \end{smallmatrix}\right) = \left(\begin{smallmatrix} gx_1 & gx_2 \end{smallmatrix}\right)$.

$$\begin{tikzcd}
X_1 \arrow[r, tail, "x_1"] \arrow[d, "{\left(\begin{smallmatrix}1\\0\end{smallmatrix}\right)}"] \arrow[dd, bend right, "f"'] & W \arrow[r, two heads] \arrow[d, equal] & Y \arrow[d] \\
X_1 \oplus X_2 \arrow[r, tail, "{\left(\begin{smallmatrix}x_1 & x_2\end{smallmatrix}\right)}"] \arrow[d, "{\left(\begin{smallmatrix}f & 0\end{smallmatrix}\right)}"] & W \arrow[r, two heads] \arrow[dl, dotted, "g"] & U \\
V_0 & &
\end{tikzcd}$$

Hence $\operatorname{Hom}_{\mathcal{B}}(x_1, V_0) : \operatorname{Hom}_{\mathcal{B}}(W, V_0) \to \operatorname{Hom}_{\mathcal{B}}(X_1, V_0)$ is surjective. By the following exact sequence

$$\operatorname{Hom}_{\mathcal{B}}(W, V_0) \xrightarrow{\operatorname{Hom}_{\mathcal{B}}(x_1, V_0)} \operatorname{Hom}_{\mathcal{B}}(X_1, V_0) \xrightarrow{0} \operatorname{Ext}^1_{\mathcal{B}}(Y, V_0) \to \operatorname{Ext}^1_{\mathcal{B}}(W, V_0) = 0$$

we have $\operatorname{Ext}^1_{\mathcal{B}}(Y, V_0) = 0$, which implies $Y \in \mathcal{U}$. □

Lemma 2.10. (a) *If* $A \overset{f}{\rightarrowtail} B \overset{g}{\twoheadrightarrow} U$ *is a short exact sequence in* $\mathcal{B}$ *with* $U \in \mathcal{U}$*, then* $A \in \mathcal{B}^-$ *implies* $B \in \mathcal{B}^-$.

(b) *If* $A \overset{f}{\rightarrowtail} B \overset{g}{\twoheadrightarrow} S$ *is a short exact sequence in* $\mathcal{B}$ *with* $S \in \mathcal{S}$*, then* $B \in \mathcal{B}^-$ *implies* $A \in \mathcal{B}^-$.

Proof. (*b*) Since $B \in \mathcal{B}^-$, by definition, there exists a short exact sequence

$$B \overset{w^B}{\rightarrowtail} W^B \twoheadrightarrow S^B.$$

Take a push-out of g and w^B, by Proposition 2.2, we get a commutative diagram of short exact sequences

$$\begin{tikzcd}
A \arrow[r, tail, "f"] \arrow[d, equal] & B \arrow[r, two heads, "g"] \arrow[d, tail, "w^B"] & S \arrow[d, tail] \\
A \arrow[r, tail] & W^B \arrow[r, two heads] \arrow[d, two heads] & X \arrow[d, two heads] \\
 & S^B \arrow[r, equal] & S^B.
\end{tikzcd}$$

We thus get $X \in \mathcal{S}$ since $\mathcal{S}$ is closed under extension. This gives $A \in \mathcal{B}^-$.
(*a*) Since $A \in \mathcal{B}^-$, it admits a short exact sequence

$$A \overset{w^A}{\rightarrowtail} W^A \twoheadrightarrow S^A.$$

where $W^A \in \mathcal{W}$ and $S^A \in \mathcal{S}$. Since $\operatorname{Ext}^1_{\mathcal{B}}(\mathcal{S}, \mathcal{T}) = 0$, w^A is a left $\mathcal{T}$-approximation of A. Thus there exists a commutative diagram of two short exact sequences

$$\begin{tikzcd}
A \arrow[r, tail] \arrow[d, "f"'] & W^A \arrow[r, two heads] \arrow[d] & S^A \arrow[d] \\
B \arrow[r, tail, "t^B"'] & T^B \arrow[r, two heads] & S^B.
\end{tikzcd}$$

It suffices to show $T^B \in \mathcal{U}$.
Apply $\operatorname{Ext}^1_{\mathcal{B}}(-, \mathcal{V})$ to the following commutative diagram

$$\begin{tikzcd}
A \arrow[r, tail, "f"] \arrow[d, "w^A"'] & B \arrow[r, two heads] \arrow[d, "t^B"] & U \\
W^A \arrow[r] & T^B &
\end{tikzcd}$$

since $\mathrm{Ext}^1_{\mathcal{B}}(\mathcal{U},\mathcal{V})=0$, we obtain the following commutative diagram

$$\begin{tikzcd}
 & \mathrm{Ext}^1_{\mathcal{B}}(T^B,\mathcal{V}) \arrow[r] \arrow[d, "\mathrm{Ext}^1_{\mathcal{B}}(t^B,\mathcal{V})"] \arrow[dl, dotted] & \mathrm{Ext}^1_{\mathcal{B}}(W^A,\mathcal{V})=0 \arrow[d] \\
0=\mathrm{Ext}^1_{\mathcal{B}}(U,\mathcal{V}) \arrow[r] & \mathrm{Ext}^1_{\mathcal{B}}(B,\mathcal{V}) \arrow[r, "\mathrm{Ext}^1_{\mathcal{B}}(f,\mathcal{V})"'] & \mathrm{Ext}^1_{\mathcal{B}}(A,\mathcal{V}).
\end{tikzcd}$$

It follows that $\mathrm{Ext}^1_{\mathcal{B}}(t^B,\mathcal{V})=0$. Then from the following exact sequence

$$0=\mathrm{Ext}^1_{\mathcal{B}}(S^B,\mathcal{V})\to\mathrm{Ext}^1_{\mathcal{B}}(T^B,\mathcal{V})\xrightarrow{\mathrm{Ext}^1_{\mathcal{B}}(t^B,\mathcal{V})=0}\mathrm{Ext}^1_{\mathcal{B}}(B,\mathcal{V})$$

we get that $\mathrm{Ext}^1_{\mathcal{B}}(T^B,\mathcal{V})=0$, which implies that $T^B\in\mathcal{U}$. Thus $T^B\in\mathcal{W}$ and $B\in B^-$. □

Dually, the following holds.

Lemma 2.11. (a) *If* $T \rightarrowtail A \overset{f}{\twoheadrightarrow} B$ *is a short exact sequence in* $\mathcal{B}$ *with* $T\in\mathcal{T}$*, then* $B\in\mathcal{B}^+$ *implies* $A\in\mathcal{B}^+$*.*

(b) *If* $V \rightarrowtail A \overset{f}{\twoheadrightarrow} B$ *is a short exact sequence in* $\mathcal{B}$ *with* $V\in\mathcal{V}$*, then* $A\in\mathcal{B}^+$ *implies* $B\in\mathcal{B}^+$*.*

Now we give a proposition which is similar with [1, Proposition 1.10] and useful in our article.

Proposition 2.12. *Let* $\mathcal{T}$ *be a subcategory of* $\mathcal{B}$ *satisfying*

(a) $\mathcal{P}\subseteq\mathcal{T}$.
(b) $\mathcal{T}$ *is contravariantly finite.*
(c) $\mathcal{T}$ *is closed under extension.*

Then we get a cotorsion pair $(\mathcal{T},\mathcal{V})$ *where*

$$\mathcal{V}=\{X\in\mathcal{B}\mid \mathrm{Ext}^1_{\mathcal{B}}(\mathcal{T},X)=0\}.$$

Proof. For any object $B\in\mathcal{B}$, it admits a short exact category

$$B \rightarrowtail I \overset{f}{\twoheadrightarrow} X$$

where $I\in\mathcal{I}$. By (a) and (b), we can take a short exact sequences

$$V_X \rightarrowtail T_X \overset{t_X}{\twoheadrightarrow} X, \qquad V_B \rightarrowtail T_B \overset{t_B}{\twoheadrightarrow} B$$

where t_X (resp. t_B) is a minimal right $\mathcal{T}$-approximation of X (resp. B). Since $\mathcal{T}$ is closed under extension, by Wakamatsu's Lemma, we obtain $V_X\in\mathcal{V}$ (resp. $V_B\in\mathcal{V}$). Take a pull-back of f and t_X, we get the following commutative diagram

$$\begin{tikzcd}
 & V_X \arrow[r, equal] \arrow[d, tail] & V_X \arrow[d, tail] \\
B \arrow[r, tail] \arrow[d, equal] & Y \arrow[r, two heads] \arrow[d] & T_X \arrow[d, "t_X"] \\
B \arrow[r, tail] & I \arrow[r, two heads, "f"'] & X.
\end{tikzcd}$$

Since $I,V\in\mathcal{V}$ and $\mathcal{V}$ is extension closed, we get $Y\in\mathcal{V}$. Thus B admits two short exact sequence

$$V_B\rightarrowtail T_B\twoheadrightarrow B,\quad B\rightarrowtail Y\twoheadrightarrow T_X$$

satisfying $V_B,Y\in\mathcal{V}$ and $T_B,T_X\in\mathcal{T}$. Hence by definition $(\mathcal{T},\mathcal{V})$ is a cotorsion pair. □

3. $\underline{\mathcal{H}}$ IS PREABELIAN

In this section, we fix a twin cotorsion pair $(\mathcal{S},\mathcal{T})$, $(\mathcal{U},\mathcal{V})$, we will show that the heart $\underline{\mathcal{H}}$ of a twin cotorsion pair is preabelian.

Definition 3.1. For any $B \in \mathcal{B}$, define B^+ and $b^+ : B \to B^+$ as follows:
Take two short exact sequences:

$$V_B \rightarrowtail U_B \twoheadrightarrow B, \quad U_B \rightarrowtail T^U \twoheadrightarrow S^U$$

where $U_B \in \mathcal{U}$, $\mathcal{V}_B \in \mathcal{V}$, $T^U \in \mathcal{T}$ and $S^U \in \mathcal{S}$. By Proposition 2.2, we get the following commutative diagram

$$\begin{tikzcd}
V_B \arrow[r, tail] \arrow[d, equal] & U_B \arrow[r, two heads] \arrow[d, tail, "u"] & B \arrow[d, tail, "b^+"] \\
V_B \arrow[r, tail] & T^U \arrow[r, two heads, "t"] \arrow[d, two heads] & B^+ \arrow[d, two heads] \\
 & S^U \arrow[r, equal] & S^U
\end{tikzcd} \tag{1}$$

where the upper-right square is both a push-out and a pull-back.

We can easily get the following Lemma.

Lemma 3.2. *By Definition 3.1, $B^+ \in \mathcal{B}^+$. Moreover, if $B \in \mathcal{B}^-$, then $B^+ \in \mathcal{H}$.*

Proof. Since $\mathcal{U}$ is closed under extension, we get $T^U \in \mathcal{U} \cap \mathcal{T} = \mathcal{W}$. Hence by definition $B^+ \in \mathcal{B}^+$. If $B \in \mathcal{B}^-$, by Lemma 2.10, B^+ also lies in $\mathcal{B}^-$. Thus $B^- \in \mathcal{H}$. $\square$

We give an important property of b^+ in the following proposition.

Proposition 3.3. *For any $B \in \mathcal{B}$ and $Y \in \mathcal{B}^+$, $\mathrm{Hom}_{\mathcal{B}}(b^+, Y) : \mathrm{Hom}_{\mathcal{B}}(B^+, Y) \to \mathrm{Hom}_{\mathcal{B}}(B, Y)$ is surjective and $\mathrm{Hom}_{\underline{\mathcal{B}}}(\underline{b^+}, Y) : \mathrm{Hom}_{\underline{\mathcal{B}}}(B^+, Y) \to \mathrm{Hom}_{\underline{\mathcal{B}}}(B, Y)$ is bijective.*

Proof. Let $y \in \mathrm{Hom}_{\mathcal{B}}(B, Y)$ be any morphism. By definition, there exists a short exact sequence

$$\begin{tikzcd}
V_Y \arrow[r, tail] & W_Y \arrow[r, two heads, "w_Y"] & Y.
\end{tikzcd}$$

Since $\mathrm{Ext}^1_{\mathcal{B}}(U_B, V_Y) = 0$, w_Y is a right $\mathcal{U}$-approximation of Y. Thus any $f \in \mathrm{Hom}_{\mathcal{B}}(U_B, Y)$ factors through W_Y.

$$\begin{tikzcd}
 & & U \arrow[dl, dotted, "g"'] \arrow[d, "f"] \\
V_Y \arrow[r, tail] & W_Y \arrow[r, two heads, "w_Y"'] & Y
\end{tikzcd}$$

As $\mathrm{Ext}^1_{\mathcal{B}}(\mathcal{S}, \mathcal{T}) = 0$, u is a left $\mathcal{T}$-approximation of U_B, we get the following commutative diagram:

$$\begin{tikzcd}
U_B \arrow[r, tail, "u"] \arrow[d, "g"] & T^U \arrow[r, two heads] \arrow[dl, dotted] & S^U \\
W_Y \arrow[d, "w_Y"] & & \\
Y & &
\end{tikzcd}$$

which implies that $\operatorname{Hom}_{\mathcal{B}}(u, Y) : \operatorname{Hom}_{\mathcal{B}}(T^U, Y) \to \operatorname{Hom}_{\mathcal{B}}(U_B, Y)$ is epimorphic. Hence when we apply $\operatorname{Hom}_{\mathcal{B}}(-, Y)$ to the diagram (3.1), we obtain the following exact sequence

$$\begin{tikzcd}
\operatorname{Hom}_{\mathcal{B}}(B^+, Y) \arrow[r, "\operatorname{Hom}_{\mathcal{B}}(b^+, Y)"] & \operatorname{Hom}_{\mathcal{B}}(B, Y) \arrow[r] \arrow[d, dotted, tail] & \operatorname{Ext}^1_{\mathcal{B}}(S, Y) \arrow[r] \arrow[d, equal] & \operatorname{Ext}^1_{\mathcal{B}}(B^+, Y) \arrow[d] \\
\operatorname{Hom}_{\mathcal{B}}(T^U, Y) \arrow[r, "\operatorname{Hom}_{\mathcal{B}}(u, Y)"] & \operatorname{Hom}_{\mathcal{B}}(U_B, Y) \arrow[r, "0"] & \operatorname{Ext}^1_{\mathcal{B}}(S^U, Y) \arrow[r] & \operatorname{Ext}^1_{\mathcal{B}}(T^U, Y)
\end{tikzcd}$$

which implies that $\operatorname{Hom}_{\mathcal{B}}(b^+, Y)$ is an epimorphism. In particular, $\operatorname{Hom}_{\underline{\mathcal{B}}}(\underline{b^+}, Y)$ is an epimorphism.
It remains to show that $\operatorname{Hom}_{\underline{\mathcal{B}}}(\underline{b^+}, Y)$ is monomorphic. Suppose $q \in \operatorname{Hom}_{\mathcal{B}}(B^+, Y)$ satisfies $\underline{qb^+} = 0$, it follows that qb^+ factors through $\mathcal{W}$. Since w_Y is a right $\mathcal{U}$-approximation, there exists a morphism $a : B \to W_Y$ such that $w_Y a = qb^+$. Take a push-out of b^+ and a, we get the following commutative diagram of short exact sequences

$$\begin{tikzcd}
B \arrow[r, tail, "b^+"] \arrow[d, "a"'] \arrow[dr, phantom, "PO"] & B^+ \arrow[r, two heads] \arrow[d, "c'"] & S^U \arrow[d, equal] \\
W_Y \arrow[r, tail, "c"'] & Q \arrow[r, two heads] & S^U.
\end{tikzcd}$$

There exists a morphism $d : Q \to Y$ such that $dc = w_Y$ and $dc' = q$ by the definition of push-out. But $Q \in \mathcal{U}$ by Lemma 2.4, and w_Y is a right $\mathcal{U}$-approximation, we have that d factors through W_Y. Thus $q = dc'$ also factors through W_Y, and $\underline{q} = 0$. ☐

We give an equivalent condition for a special case when $B^+ = 0$ in $\underline{\mathcal{B}}$.

Lemma 3.4. *For any $B \in \mathcal{B}$, the following are equivalent.*

(a) $B^+ \in \mathcal{W}$.
(b) $B \in \mathcal{U}$.
(c) $\underline{b^+} = 0$ *in* $\underline{\mathcal{B}}$.

Proof. Consider the diagram (3.1) in Definition 3.1. We first prove that $(a) \Leftrightarrow (b)$.
Suppose (b) holds. Since $B \in \mathcal{U}$, we get $B^+ \in \mathcal{U}$. Thus $\operatorname{Ext}^1_{\mathcal{B}}(B^+, V_B) = 0$, and then t splits. Hence B^+ is a direct summand of $T^U \in \mathcal{W}$, which implies that $B^+ \in \mathcal{W}$.
Now Suppose (a) holds, then the second row splits. Thus we get a pull-back diagram

$$\begin{tikzcd}
U_B \arrow[r, two heads] \arrow[d, tail] & B \arrow[d, tail, "b^+"] \\
V_B \oplus B^+ \arrow[r, two heads, "(0\ 1)"'] & B^+
\end{tikzcd}$$

which implies that u also splits, hence $B \in \mathcal{U}$.
Obviously (a) implies (c), now it suffices to show that (c) implies (b).
Since b^+ factors through $\mathcal{W}$, and t is a right $\mathcal{U}$-approximation of B^+, we get that b^+ factors through t. Hence by the definition of pull-back, the first row of diagram (3.1) splits, which implies that $B \in \mathcal{U}$. ☐

Now we give a dual construction.

Definition 3.5. For any object $B \in \mathcal{B}$, we define $b^- : B^- \to B$ as follows Take the following two short exact sequences

$$B \rightarrowtail T^B \twoheadrightarrow S^B, \quad V_T \rightarrowtail U_T \twoheadrightarrow T^B$$

where $U_T \in \mathcal{U}$, $V_T \in \mathcal{V}$, $T^B \in \mathcal{T}$ and $S^B \in \mathcal{S}$. By Proposition 2.2, we get the following commutative diagram:

$$\begin{tikzcd}
V_T \arrow[r, equal] \arrow[d, tail] & V_T \arrow[d, tail] & \\
B^- \arrow[r, tail] \arrow[d, two heads, "b^-"'] & U_T \arrow[r, two heads] \arrow[d, two heads] & S^B \arrow[d, equal] \\
B \arrow[r, tail] & T^B \arrow[r, two heads] & S^B.
\end{tikzcd}$$

By duality, we get:

Proposition 3.6. *For any* $B \in \mathcal{B}$, $B^- \in \mathcal{B}^-$ *and* $B \in \mathcal{B}^+$ *implies* $B^- \in \mathcal{H}$. *For any* $X \in \mathcal{B}^-$, $\operatorname{Hom}_{\mathcal{B}}(X, b^-) : \operatorname{Hom}_{\mathcal{B}}(X, B^-) \to \operatorname{Hom}_{\mathcal{B}}(X, B)$ *is surjective and* $\operatorname{Hom}_{\underline{\mathcal{B}}}(X, \underline{b^-}) : \operatorname{Hom}_{\underline{\mathcal{B}}}(X, B^-) \to \operatorname{Hom}_{\underline{\mathcal{B}}}(X, B)$ *is bijective.*

Definition 3.7. For any morphism $f : A \to B$ with $A \in \mathcal{B}^-$, define C_f and $c_f : B \to C_f$ as follows: By definition, there exists a short exact sequence

$$\begin{tikzcd}
A \arrow[r, tail, "w^A"] & W^A \arrow[r, two heads] & S^A.
\end{tikzcd}$$

Take a push-out of f and w^A, we get the following commutative diagram of short exact sequences

$$\begin{tikzcd}
A \arrow[r, tail, "w^A"] \arrow[d, "f"'] \arrow[dr, phantom, "PO"] & W^A \arrow[r, two heads] \arrow[d] & S^A \arrow[d, equal] \\
B \arrow[r, tail, "c_f"'] & C_f \arrow[r, two heads, "s"'] & S^A.
\end{tikzcd}\tag{2}$$

By Lemma 2.10, $B \in \mathcal{B}^-$ implies $C_f \in \mathcal{B}^-$.

Dually, we have the following:

Definition 3.8. For any morphism $f : A \to B$ in $\mathcal{B}$ with $B \in \mathcal{B}^+$, define K_f and $k_f : K_f \to A$ as follows: By definition, there exists a short exact sequence

$$\begin{tikzcd}
V_B \arrow[r, tail] & W_B \arrow[r, two heads, "w_B"] & B.
\end{tikzcd}$$

Take a pull-back of f and w_B, we get the following commutative diagram of short exact sequences

$$\begin{tikzcd}
V_B \arrow[r, tail] \arrow[d, equal] & K_f \arrow[r, two heads, "k_f"] \arrow[d] \arrow[dr, phantom, "PB"] & A \arrow[d, "f"] \\
V_B \arrow[r, tail] & W_B \arrow[r, two heads, "w_B"'] & B
\end{tikzcd}\tag{3}$$

By Lemma 2.11, $A \in B^+$ implies $K_f \in B^+$.

The following lemma gives an important property of c_f:

Lemma 3.9. *Let* $f : A \to B$ *be any morphism in* $\mathcal{B}$ *with* $A \in \mathcal{B}^-$, *take the notation of Definition 3.7, then* $c_f : B \to C_f$ *satisfying the following properties:*

For any $C \in \mathcal{B}$ and any morphism $g \in \mathrm{Hom}_{\mathcal{B}}(B, C)$ satisfying $\underline{gf} = 0$, there exists a morphism $c : C_f \to C$ such that $cc_f = g$.

$$\begin{tikzcd}
A \arrow[r, "f"] & B \arrow[rr, "g"] \arrow[dr, "c_f"'] & & C \\
& & C_f \arrow[ur, dotted, "c"'] &
\end{tikzcd}$$

Moreover if $C \in \mathcal{B}^+$, then $\underline{c}$ is unique in $\underline{\mathcal{B}}$. The dual statement also holds for k_f in Definition 3.8.

Proof. Since $\underline{gf} = 0$, gf factors through $\mathcal{W}$. As $\mathrm{Ext}^1_{\mathcal{B}}(S_A, W) = 0$, w^A is a left $\mathcal{W}$-approximation of A. Hence there exists $b : W^A \to C$ such that $gf = bw^A$. Then by the definition of push-out, we get the following commutative diagram

$$\begin{tikzcd}
A \arrow[r, "w^A"] \arrow[d, "f"'] & W^A \arrow[d] \arrow[ddr, bend left, "b"] & \\
B \arrow[r, "c_f"] \arrow[drr, bend right, "g"'] & C_f \arrow[dr, dotted, "c"] & \\
& & C.
\end{tikzcd}$$

Now assume that $C \in \mathcal{B}^+$ and there exists $c' : C_f \to C$ such that $c'c_f = g$. Since $(c' - c)c_f = 0$, there exists a morphism $d : S^A \to C$ such that $c' - c = ds$. As C admits a short exact sequence

$$\begin{tikzcd}
V_C \arrow[r, tail] & W_C \arrow[r, two heads, "w_C"] & C
\end{tikzcd}$$

and w_C is a right $\mathcal{U}$-approximation of C, we obtain that there exists a morphism $e : S^A \to W_C$ such that $w_C e = d$. Hence $c' - c$ factors through W_C, and $\underline{c} = \underline{c'}$. □

Theorem 3.10. *For any twin cotorsion pair $(\mathcal{S}, \mathcal{T}), (\mathcal{U}, \mathcal{V})$, its heart $\underline{\mathcal{H}}$ is preabelian.*

Proof. We only show the construction of the cokernel. For any $A, B \in \mathcal{H}$ and any morphism $f : A \to B$, by Definition 3.7, since $A, B \in \mathcal{B}^-$, it follows $\underline{c_f f} = 0$ and $C_f \in \mathcal{B}^-$. By Proposition 3.3, there exists ${c_f}^+ : C_f \to C_f$ where ${C_f}^+ \in \mathcal{H}$ by Lemma 3.2. We claim that $\underline{{c_f}^+ c_f} : B \to {C_f}^+$ is the cokernel of $\underline{f}$. Let Q be any object in $\mathcal{H}$, and let $r : B \to Q$ be any morphism satisfying $\underline{rf} = 0$, then by Proposition 3.9 and 3.3, there exists a commutative diagram

$$\begin{tikzcd}
& Q & & \\
A \arrow[ur, "0"] \arrow[r, "\underline{f}"'] & B \arrow[u, "\underline{r}"] \arrow[r, "\underline{c_f}"'] & C_f \arrow[ul, dotted, "\underline{a}"] \arrow[r, "\underline{{c_f}^+}"'] & {C_f}^+. \arrow[ull, dotted, "\underline{b}"']
\end{tikzcd}$$

The uniqueness of $\underline{b}$ follows from Lemma 3.9 and Proposition 3.3. □

Corollary 3.11. *Let $f : A \to B$ be a morphism in $\mathcal{H}$, the the followings are equivalent:*

(a) *$\underline{f}$ is epimorphic in $\underline{\mathcal{H}}$.*
(b) *${C_f}^+ \in \mathcal{W}$.*
(c) *$C_f \in \mathcal{U}$.*

Proof. The equivalence of (b) and (c) is given by Lemma 3.4. We prove the equivalence of (a) and (b). By Theorem 3.10, $\underline{{c_f}^+ c_f}$ is the cokernel of $\underline{f}$ in $\underline{\mathcal{H}}$. Then (b) implies that $\underline{{c_f}^+ c_f} = 0$ in $\underline{\mathcal{H}}$, which means $\underline{f}$ is epimorphic in $\underline{\mathcal{H}}$.
If (a) holds, then $\underline{{c_f}^+ c_f} = 0$ since the cokernel of $\underline{f}$ in $\underline{\mathcal{H}}$ is 0. By Lemma 3.9, we get $\underline{{c_f}^+} = 0$. By Lemma 3.4, ${C_f}^+ \in \mathcal{W}$. □

4. ABELIANESS OF THE HEARTS OF COTORSION PAIRS

In this section we fix a cotorsion pair $(\mathcal{U},\mathcal{V})$. We will prove that the heart $\underline{\mathcal{H}}=\mathcal{B}^+\cap\mathcal{B}^-/\mathcal{U}\cap\mathcal{V}$ of a cotorsion pair is abelian.

Lemma 4.1. *Let $A,B\in\mathcal{H}$, and let*

$$C\rightarrowtail^{g} A \xrightarrow{f}\!\!\!\!\rightarrow B$$

be a short exact sequence in $\mathcal{B}$. If $\underline{f}$ is epimorphic in $\underline{\mathcal{H}}$, then C belongs to $\mathcal{B}^-$.

Proof. As $\underline{f}$ is epimorphic in $\underline{\mathcal{H}}$, we get $C_f\in\mathcal{U}$ by Corollary 5.3. By Definition 3.7, we get following commutative diagram

$$\begin{tikzcd}
C \arrow[r, equal] \arrow[d, tail, "g"'] & C \arrow[d, tail, "h"] & \\
A \arrow[r, tail, "w^A"] \arrow[d, "f"'] \arrow[rd, phantom, "PO"] & W^A \arrow[r, two heads] \arrow[d] & S^A \arrow[d, equal] \\
B \arrow[r, tail, "c_f"'] & C_f \arrow[r, two heads] & S^A.
\end{tikzcd} \tag{4}$$

The middle column shows that $C\in\mathcal{B}^-$. □

We need the following lemma to prove our theorem.

Lemma 4.2. (a) *Let $f:A\to B$ be a morphism in $\mathcal{B}$ with $B\in\mathcal{B}^+$, then there exists a deflation $\alpha=(\,f\;-w_B\,):A\oplus W_B\twoheadrightarrow B$ in $\mathcal{B}$ such that $\underline{\alpha}=\underline{f}$.*

(b) *Let $f:A\to B$ be a morphism in $\mathcal{B}$ with $A\in\mathcal{B}^-$, then there exists an inflation $\alpha=\begin{pmatrix} f\\ -w^A\end{pmatrix}$: $A\rightarrowtail B\oplus W^A$ in $\mathcal{B}$ such that $\underline{\alpha'}=\underline{f}$.*

Proof. We only show the first one, the second is dual.
As $B\in\mathcal{B}^+$, it admits a short exact sequence

$$V_B\rightarrowtail W_B \xrightarrow{w_B}\!\!\!\!\rightarrow B.$$

Take a pull-back of f and w_B, by dual of Proposition 2.1, we get a short exact sequence

$$C\rightarrowtail A\oplus W_B \xrightarrow{\alpha=(\,f\;-w_B\,)}\!\!\!\!\rightarrow B$$

and consequently α is a deflation and $\underline{\alpha}=\underline{f}$. □

Theorem 4.3. *For any cotorsion pair $(\mathcal{U},\mathcal{V})$ on $\mathcal{B}$, its heart $\underline{\mathcal{H}}$ is an abelian category.*

Proof. Since $\mathcal{H}$ is preabelian, it remains to show the following:

(a) If $\underline{f}$ is epimorphic in $\underline{\mathcal{H}}$, then $\underline{f}$ is a cokernel of some morphism in $\underline{\mathcal{H}}$.
(b) If $\underline{f}$ is monomorphic in $\underline{\mathcal{H}}$, then $\underline{f}$ is a kernel of some morphism in $\underline{\mathcal{H}}$.

We only show (a), since (b) is dual.
For any morphism $\underline{f}:A\to B$ which is epimorphic in $\underline{\mathcal{H}}$, by Lemma 4.2, it is enough to consider the case that f is a deflation.
Let f admit a short exact sequence:

$$C\rightarrowtail^{g} A \xrightarrow{f}\!\!\!\!\rightarrow B.$$

By Lemma 4.1, we have $C\in\mathcal{B}^-$. By Proposition 3.3, there exists

$$c^+:C\to C^+$$

where C^+ lies in $\mathcal{H}$ by Lemma 3.2. As $A \in \mathcal{B}^+$, there exists $a : C^+ \to A$ such that $ac^+ = g$.

$$\begin{tikzcd}
C \arrow[rr, "g"] \arrow[dr, "c^+"'] & & A \\
& C^+. \arrow[ur, "a"'] &
\end{tikzcd}$$

Since $\underline{fac^+} = \underline{fg} = 0$, we have $\underline{fa} = 0$ by Proposition 3.3. We claim that $\underline{f}$ is the cokernel of $\underline{a}$.
Let Q be any object in $\mathcal{H}$ and $r : A \to Q$ be any morphism. By Proposition 3.3, $\underline{rg} = 0$ if and only if $\underline{ra} = 0$.
So it is enough to show that any $\underline{r}$ satisfying $\underline{rg} = 0$ factors through $\underline{f}$.
If $\underline{rg} = 0$, rg factors through $\mathcal{W}$. Consider the second column of diagram (4), since h is a left $\mathcal{V}$-approximation of C, there exists a morphism $c : W^A \to Q$ such that $rg = ch$. Since $h = w^A g$, we get that $(r - w^A)g = 0$. Thus $r - w^A$ factors through f, which implies that $\underline{r}$ factors through $\underline{f}$. □

5. $\underline{\mathcal{H}}$ IS SEMI-ABELIAN

In the following sections, we fix a twin cotorsion pair $(\mathcal{S}, \mathcal{T}), (\mathcal{U}, \mathcal{V})$.

Definition 5.1. A preabelian category $\mathcal{A}$ is called *left semi-abelian* if in any pull-back diagram

$$\begin{tikzcd}
A \arrow[r, "\alpha"] \arrow[d, "\beta"'] & B \arrow[d, "\gamma"] \\
C \arrow[r, "\delta"'] & D
\end{tikzcd}$$

in $\mathcal{A}$, α is an epimorphism whenever δ is a cokernel. *Right semi-abelian* is defined dually. $\mathcal{A}$ is called *semi-abelian* if it is both left and right semi-abelian. In this section we will prove that the heart $\underline{\mathcal{H}}$ of a twin cotorsion pair is semi-ableian.

Lemma 5.2. *If morphism $\beta \in \mathrm{Hom}_{\underline{\mathcal{H}}}(B, C)$ is a cokernel of a morphism $\underline{f} \in \mathrm{Hom}_{\underline{\mathcal{H}}}(A, B)$, then B admits a short exact sequence*

$$B \rightarrowtail C' \twoheadrightarrow S$$

where $C' \in \mathcal{H}$, $C \simeq C'$ in $\underline{\mathcal{H}}$ and $S \in \mathcal{S}$.

Proof. Let β be the cokernel of $f : A \to B$. By Theorem 3.10, the cokernel of $\underline{f}$ is given by $\underline{{c_f}^+ c_f}$. Therefore ${C_f}^+ \simeq C$ in $\underline{\mathcal{H}}$. Consider diagram (4) and the diagram which induces $(C_f)^+$ by Definition 3.1:

$$\begin{tikzcd}
V \arrow[r, tail] \arrow[d, equal] & U \arrow[r, two heads] \arrow[d, tail] & C_f \arrow[d, tail, "{c_f}^+"] \\
V \arrow[r, tail] & T' \arrow[r, two heads] \arrow[d, two heads] & {C_f}^+ \arrow[d, two heads] \\
& S' \arrow[r, equal] & S'
\end{tikzcd}$$

by Proposition 2.2, we obtain the following commutative diagram of short exact sequences

$$\begin{tikzcd}
B \arrow[r, tail, "c_f"] \arrow[d, equal] & C_f \arrow[r, two heads] \arrow[d, tail, "{c_f}^+"'] & S^A \arrow[d, tail] \\
B \arrow[r, tail] & {C_f}^+ \arrow[r, two heads] \arrow[d, two heads] & Q \arrow[d, two heads] \\
& S' \arrow[r, equal] & S'.
\end{tikzcd}$$

From the third column we get $Q \in \mathcal{S}$. Hence we get the required short exact sequence. $\square$

Proposition 5.3. *Let* $A \overset{f}{\rightarrowtail} B \overset{g}{\twoheadrightarrow} C$ *be a short exact sequence in* $\mathcal{B}$ *with* f *in* $\mathcal{H}$*. If* g *factors through* $\mathcal{U}$*, then* $\underline{f}$ *is epimorphic in* $\underline{\mathcal{H}}$*.*

Proof. By Corollary 3.11, it suffices to show that $C_f \in \mathcal{U}$.
By definition of $c_f : B \to C_f$, there is a commutative diagram of short exact sequences

$$\begin{tikzcd}
A \arrow[r, tail, "w^A"] \arrow[d, tail, "f"'] \arrow[dr, phantom, "PO"] & W^A \arrow[r, two heads] \arrow[d] & S^A \arrow[d, equal] \\
B \arrow[r, tail, "c_f"'] \arrow[d, two heads, "g"'] & C_f \arrow[r, two heads] \arrow[d, two heads] & S^A \\
C \arrow[r, equal] & C. &
\end{tikzcd}$$

Since $\operatorname{Ext}^1_{\mathcal{B}}(\mathcal{W}, \mathcal{V}) = 0$, we get the following commutative diagram of exact sequence

$$\begin{tikzcd}
\operatorname{Ext}^1_{\mathcal{B}}(C,\mathcal{V}) \arrow[r] \arrow[d, equal] & \operatorname{Ext}^1_{\mathcal{B}}(C_f,\mathcal{V}) \arrow[r] \arrow[d, "\operatorname{Ext}^1_{\mathcal{B}}(c_f{,}\mathcal{V})"] & \operatorname{Ext}^1_{\mathcal{B}}(W^A,\mathcal{V}) = 0 \arrow[d] \\
\operatorname{Ext}^1_{\mathcal{B}}(C,\mathcal{V}) \arrow[r, "\operatorname{Ext}^1_{\mathcal{B}}(g{,}\mathcal{V})"'] & \operatorname{Ext}^1_{\mathcal{B}}(B,\mathcal{V}) \arrow[r, "\operatorname{Ext}^1_{\mathcal{B}}(f{,}\mathcal{V})"'] & \operatorname{Ext}^1_{\mathcal{B}}(A,\mathcal{V}).
\end{tikzcd}$$

Then $\operatorname{Ext}^1_{\mathcal{B}}(c_f, \mathcal{V})$ factors through $\operatorname{Ext}^1_{B}(g, \mathcal{V})$. We have $\operatorname{Ext}^1_{\mathcal{B}}(g, \mathcal{V}) = 0$ since g factors through $\mathcal{U}$, thus we get $\operatorname{Ext}^1_{\mathcal{B}}(f, \mathcal{V}) = 0$. Then from the following exact sequence

$$0 = \operatorname{Ext}^1_{\mathcal{B}}(S^A, \mathcal{V}) \to \operatorname{Ext}^1_{\mathcal{B}}(C_f, \mathcal{V}) \xrightarrow{\operatorname{Ext}^1_{\mathcal{B}}(f,\mathcal{V})=0} \operatorname{Ext}^1_{\mathcal{B}}(B, \mathcal{V})$$

we obtain that $\operatorname{Ext}^1_{\mathcal{B}}(C_f, \mathcal{V}) = 0$, which implies $C_f \in \mathcal{U}$.

$\square$

Lemma 5.4. *Suppose* $X \in \mathcal{B}^-$ *admits a short exact sequence*

$$X \overset{x}{\rightarrowtail} B \twoheadrightarrow U$$

where $B \in \mathcal{H}$ *and* $U \in \mathcal{U}$*. Then the unique morphism* $\underline{b} \in \operatorname{Hom}_{\underline{\mathcal{H}}}(X^+, B)$ *given by Proposition 3.3 which satisfies* $\underline{bx^+} = \underline{x}$ *is epimorphic.*

Proof. By Definition 3.1, there exists a short exact sequence

$$X \overset{x^+}{\rightarrowtail} X^+ \twoheadrightarrow S$$

where $S \in \mathcal{S}$. By Proposition 3.3, there exits $b : X^+ \to B$ such that $bx^+ = x$. Since $X \in \mathcal{B}^-$, we obtain $C^+ \in \mathcal{H}$ by Lemma 3.2. Hence X^+ admits a short exact sequence

$$X^+ \overset{a}{\rightarrowtail} W \twoheadrightarrow S'$$

where $W \in \mathcal{W}$ and $S' \in \mathcal{S}$. Take a push-out of a and b, we get the following commutative diagram

$$\begin{tikzcd}
X^+ \arrow[r, tail, "a"] \arrow[d, "b"'] & W \arrow[r, two heads] \arrow[d] & S' \arrow[d, equal] \\
B \arrow[r, tail] & C \arrow[r, two heads] & S'
\end{tikzcd}$$

which induces a short exact sequence

$$X^+ \overset{\binom{b}{-a}}{\rightarrowtail} B \oplus W \twoheadrightarrow C$$

by Proposition 2.1. By Proposition 2.2, we obtain the following commutative diagram

$$\begin{tikzcd}
X \arrow[r, tail, "x^+"] \arrow[d, equal] & X^+ \arrow[r, two heads] \arrow[d, tail, "a"] & S \arrow[d, tail] \\
X \arrow[r, tail, "c"'] & W \arrow[r, two heads] \arrow[d, tail] & Q \arrow[d] \\
 & S' \arrow[r, equal] & S'.
\end{tikzcd}$$

Take a push-out of x and c

$$\begin{tikzcd}
X \arrow[r, tail, "c"] \arrow[d, tail, "x"'] & W \arrow[r, two heads] \arrow[d] & Q \arrow[d, equal] \\
B \arrow[r, tail] \arrow[d] & C' \arrow[r, two heads] \arrow[d] & Q \\
U \arrow[r, equal] & U &
\end{tikzcd}$$

from the second column we obtain that $C' \in \mathcal{U}$ and we get the following short exact sequence

$$X \xrightarrow{\left(\begin{smallmatrix} x \\ -c \end{smallmatrix}\right)} B \oplus W \twoheadrightarrow C'$$

by Proposition 2.1. Thus we get the following commutative diagram

$$\begin{tikzcd}
 & X \arrow[dl, "x^+"'] \arrow[dr, tail, "\left(\begin{smallmatrix} x \\ -c \end{smallmatrix}\right)"] \arrow[d, phantom, "\circlearrowright"] & & & \\
X^+ \arrow[rr, tail, "\left(\begin{smallmatrix} b \\ -a \end{smallmatrix}\right)"'] & {} & B \oplus W \arrow[rr, two heads] \arrow[dr] & & C. \\
 & & & C' \arrow[ur, dotted] &
\end{tikzcd}$$

Hence by Proposition 5.3, $\underline{b}$ is epimorphic. ☐

We introduce the following lemma which is an analog of [14, Lemma 5.3].

Lemma 5.5. *Let*

$$\begin{tikzcd}
A \arrow[r, "\alpha"] \arrow[d, "\beta"'] & B \arrow[d, "\gamma"] \\
C \arrow[r, "\delta"'] & D
\end{tikzcd}$$

be a pull-back diagram in $\underline{\mathcal{H}}$. If there exists an object $X \in \mathcal{B}^-$ and morphisms $x_B : X \to B$, $x_C : X \to C$ which satisfy the following conditions, then α is epimorphic in $\underline{\mathcal{H}}$.

(a) *The following diagram commutative.*

$$\begin{tikzcd}
X \arrow[r, "\underline{x_B}"] \arrow[d, "\underline{x_C}"'] & B \arrow[d, "\gamma"] \\
C \arrow[r, "\delta"'] & D
\end{tikzcd}$$

(b) *There exists a short exact sequence $X \xrightarrow{x_B} B \to U$ with $U \in \mathcal{U}$.*

Proof. Take $x^+ : X \to X^+$ as in Definition 3.1. Then by Proposition 3.3, there exist $f_B : X^+ \to B$ and $f_C : X^+ \to C$ such that $\underline{f_B x^+} = \underline{x_B}$ and $\underline{f_C x^+} = \underline{x_C}$. By Lemma 5.4, $\underline{f_B}$ is epimorphic in $\underline{\mathcal{H}}$. As $\gamma \underline{x_B} = \delta \underline{x_C}$, we get $\gamma \underline{f_B x^+} = \delta \underline{f_C x^+}$, it follows by Proposition 3.3 that $\gamma \underline{f_B} = \delta \underline{f_C}$. By the definition of pull-back, there exists a morphism $\eta : X^+ \to A$ in $\underline{\mathcal{H}}$ which makes the following diagram commute.

$$\begin{tikzcd}
X^+ \arrow[rrd, bend left, "\underline{f_B}"] \arrow[rd, dotted, "\eta"] \arrow[rdd, bend right, "\underline{f_C}"'] & & \\
& A \arrow[r, "\alpha"] \arrow[d, "\beta"] & B \arrow[d, "\gamma"] \\
& C \arrow[r, "\delta"'] & D
\end{tikzcd}$$

Since $\underline{f_B}$ is epimorphic, we obtain that α is also epimorphic. □

Theorem 5.6. *For any twin cotorsion pair* $(\mathcal{S}, \mathcal{T}), (\mathcal{U}, \mathcal{V})$*, its heart* $\underline{\mathcal{H}}$ *is semi-abelian.*

Proof. By duality, we only show $\underline{\mathcal{H}}$ is left semi-abelian. Assume we are given a pull-back diagram

$$\begin{tikzcd}
A \arrow[r, "\alpha"] \arrow[d, "\beta"'] & B \arrow[d, "\gamma"] \\
C \arrow[r, "\delta"'] & D
\end{tikzcd}$$

in $\underline{\mathcal{H}}$ where δ is a cokernel. It suffices to show that α becomes epimorphic.
By Lemma 5.2, replacing D by an isomorphic one if necessary, we can assume that there exists an inflation $d : C \rightarrowtail D$ satisfying $\delta = \underline{d}$, which admits a short exact sequence

$$\begin{tikzcd}
C \arrow[r, tail, "d"] & D \arrow[r, two heads] & S
\end{tikzcd}$$

where $S \in \mathcal{S}$. As $D \in \mathcal{B}^+$, by Lemma 4.2 we can also assume that there exists an deflation $c : B \twoheadrightarrow D$ such that $\gamma = \underline{c}$. By Proposition 2.2, we get the following commutative diagram of short exact sequences

$$\begin{tikzcd}
X \arrow[r, tail, "x_B"] \arrow[d, two heads, "x_C"'] & B \arrow[r, two heads] \arrow[d, two heads, "c"] & S \arrow[d, equal] \\
C \arrow[r, tail, "d"'] & D \arrow[r, two heads] & S.
\end{tikzcd}$$

it follows by Lemma 2.10 that $X \in \mathcal{B}^-$. Hence by Lemma 5.5 α is epimorphic in $\underline{\mathcal{H}}$. □

6. The case where $\underline{\mathcal{H}}$ becomes integral

Definition 6.1. A preabelian category $\mathcal{A}$ is called *left integral* if in any pull-back diagram

$$\begin{tikzcd}
A \arrow[r, "\alpha"] \arrow[d, "\beta"'] & B \arrow[d, "\gamma"] \\
C \arrow[r, "\delta"'] & D
\end{tikzcd}$$

in $\mathcal{A}$, α is an epimorphism whenever δ is an epimorphic. *Right integral* is defined dually. $\mathcal{A}$ is called *integral* if it is both left and right integral.

In this section we give a sufficient condition where the heart $\underline{\mathcal{H}}$ becomes integral.
Let $\mathcal{C}$ be a subcategory of $\mathcal{B}$, denote by $\Omega\mathcal{C}$ (resp. $\Omega^-\mathcal{C}$) the subcateogy of $\mathcal{B}$ consisting of objects ΩC (resp. $\Omega^- C$) such that there exists a short exact sequence

$$\Omega C \rightarrowtail P_C \twoheadrightarrow C \ (P \in \mathcal{P}, C \in \mathcal{C})$$
$$(\text{resp. } C \rightarrowtail I^C \twoheadrightarrow \Omega^- C \ (I \in \mathcal{I}, C \in \mathcal{C})).$$

By definition we get $\mathcal{P} \subseteq \Omega\mathcal{C}$ and $\mathcal{I} \subseteq \Omega^{-}\mathcal{C}$. By Lemma 2.9 we get that for any cotorsion pair $(\mathcal{U}, \mathcal{V})$ on $\mathcal{B}$, $\Omega\mathcal{U}$ and $\Omega^{-}\mathcal{V}$ are closed under direct summands.

Let $\mathcal{B}_1$ $\mathcal{B}_2$ be two subcategories of $\mathcal{B}$, recall that $\mathcal{B}_1 * \mathcal{B}_2$ is subcategory of $\mathcal{B}$ consisting of objects X such that there exists a short exact sequence

$$B_1 \rightarrowtail X \twoheadrightarrow \mathcal{B}_2$$

where $B_1 \in \mathcal{B}_1$ and $B_2 \in \mathcal{B}_2$.

Theorem 6.2. *If a twin cotorsion pair* $(\mathcal{S}, \mathcal{T}), (\mathcal{U}, \mathcal{V})$ *satisfies*

$$\mathcal{U} \subseteq \mathcal{S} * \mathcal{T},\ \mathcal{P} \subseteq \mathcal{W} \quad \text{or} \quad \mathcal{T} \subseteq \mathcal{U} * \mathcal{V},\ \mathcal{I} \subseteq \mathcal{W}$$

then $\underline{\mathcal{H}}$ *becomes integral.*

Proof. According to [15, Proposition 6], a semi-abelian category is left integral if and only if it is right integral. By duality, it suffices to show that $\mathcal{U} \subseteq \mathcal{S} * \mathcal{T}, \mathcal{P} \subseteq \mathcal{W}$ implies that $\underline{\mathcal{H}}$ is left integral. Assume we are given a pull-back diagram

$$\begin{tikzcd}
A \arrow[r, "\alpha"] \arrow[d, "\beta"'] & B \arrow[d, "\gamma"] \\
C \arrow[r, "\delta"'] & D
\end{tikzcd}$$

in $\underline{\mathcal{H}}$ where δ is an epimorphism. It is sufficient to show that α is epimorphic.
Let $d : C \to D$ and $c : B \to D$ be morphisms satisfying $\delta = \underline{d}$ and $\gamma = \underline{c}$. Since δ is epimorphic, if we take $c_d : D \to C_d$ as in Definition 3.7

$$\begin{tikzcd}
C \arrow[r, tail, "w^C"] \arrow[d, "d"'] \arrow[dr, phantom, "PO"] & W^C \arrow[r, two heads] \arrow[d] & S^C \arrow[d, equal] \\
D \arrow[r, tail, "c_d"'] & C_d \arrow[r, two heads, "r"'] & S^C
\end{tikzcd}$$

then $C_d \in \mathcal{U}$ by Lemma 3.4. By assumption $\mathcal{U} \subseteq \mathcal{S} * \mathcal{T}$, C_d admits a short exact sequence

$$\begin{tikzcd}
S_0 \arrow[r, tail, "s_0"] & C_d \arrow[r, two heads, "t_0"] & T_0
\end{tikzcd}$$

with $S_0 \in \mathcal{S}$, $T_0 \in \mathcal{T}$. Since $B \in \mathcal{B}^-$ admits a short exact sequence

$$B \rightarrowtail W^B \twoheadrightarrow S^B$$

and S^B admits a short exact sequence

$$\begin{tikzcd}
\Omega S^B \arrow[r, "p"] & P_{S^B} \arrow[r, two heads, "s"] & S^B.
\end{tikzcd}$$

Thus there exists a commutative diagram

$$\begin{tikzcd}
\Omega S^B \arrow[r, tail, "p"] \arrow[d, "s_B"'] & P_{S^B} \arrow[r, two heads, "s"] \arrow[d] & S^B \arrow[d, equal] \\
B \arrow[r, tail] & W^B \arrow[r, two heads] & S^B.
\end{tikzcd} \tag{5}$$

As $\mathrm{Ext}^1_{\mathcal{B}}(S_B, T_0) = 0$, p is a left $\mathcal{T}$-approximation of ΩS^B. Therefore there exists a morphism $f : P_{S^B} \to T_0$ such that $t_0 c_d c s_B = fp$. As $P_{S^B} \in \mathcal{P}$, we conclude that there is a morphism $h : P_{S^B} \to C_d$ such that

$f=t_0h$. Since $t_0(c_dcs_B-hp)=0$, there exists a morphism $g:\Omega S_B\to S_0$ such that $c_dcs_B-hp=s_0g$. Then we get the following diagram

$$\begin{tikzcd}
\Omega S^B \arrow[rr, tail, "p"] \arrow[d, "s_B"'] \arrow[dr, dotted, "g"] & & P_{S^B} \arrow[r, two heads, "s"] \arrow[ddl, dotted, "h"'] \arrow[dddl, dotted, "f"] & S^B \\
B \arrow[d, "c"'] & S_0 \arrow[d, tail, "s_0"'] & & \\
D \arrow[r, "c_d"] & C_d \arrow[d, "t_0"'] & & \\
 & T_0. & &
\end{tikzcd}$$

Take a push-out of p and g, we get the following commutative diagram

$$\begin{tikzcd}
\Omega S^B \arrow[r, tail, "p"] \arrow[d, "g"'] & P_{S^B} \arrow[r, two heads, "s"] \arrow[d] & S^B \arrow[d, equal] \\
S_0 \arrow[r, tail] & Q \arrow[r, two heads] & S^B
\end{tikzcd}$$

and a short exact sequence

$$\begin{tikzcd}
\Omega S^B \arrow[r, tail, "\binom{p}{-g}"] & P_{S^B}\oplus S_0 \arrow[r, two heads] & Q
\end{tikzcd}$$

by Proposition 2.1 where $Q\in\mathcal{S}$. As Q admits a short exact sequence

$$\begin{tikzcd}
\Omega Q \arrow[r, tail, "k_Q"] & P_Q \arrow[r, two heads, "l_Q"] & Q
\end{tikzcd}$$

we get the following commutative diagram of short exact sequences

$$\begin{tikzcd}
\Omega Q \arrow[r, tail, "k_Q"] \arrow[d, "q_B"'] & P_Q \arrow[r, two heads, "l_Q"] \arrow[d] & Q \arrow[d, equal] \\
\Omega S^B \arrow[r, tail, "\binom{p}{-g}"] & P_{S^B}\oplus S_0 \arrow[r, two heads] & Q.
\end{tikzcd} \tag{6}$$

Since $c_dcs_B=hp+s_0g$, we obtain the following commutative diagram of short exact sequences.

$$\begin{tikzcd}
\Omega S^B \arrow[r, tail, "\binom{p}{-g}"] \arrow[d, "cs_B"'] & P_{S^B}\oplus S_0 \arrow[r, two heads] \arrow[d, "(h\ -s_0)"] & Q \arrow[d] \\
D \arrow[r, tail, "c_d"'] & C_d \arrow[r, two heads, "r"'] & S^C
\end{tikzcd}$$

Thus we get the following commutative diagram

$$\begin{tikzcd}
\Omega Q \arrow[r, tail, "k_Q"] \arrow[d, "cs_Bq_B"'] & P_Q \arrow[r, two heads, "l_Q"] \arrow[d, "n_Q"] & Q \arrow[d] \\
D \arrow[r, tail, "c_d"'] & C_d \arrow[r, two heads, "r"'] & S^C.
\end{tikzcd}$$

As $\mathcal{P} \subseteq \mathcal{W}$, we conclude that $\Omega Q \in \mathcal{B}^-$. Since S^C admits a short exact sequence

$$\Omega S^C \overset{k_{S^C}}{\rightarrowtail} P_{S^C} \overset{l_{S^C}}{\twoheadrightarrow} S^C$$

where $P_{S^C} \in \mathcal{P}$, hence we get the following commutative diagram of short exact sequence

$$\begin{tikzcd}
\Omega S^C \arrow[r, tail, "k_{S^C}"] \arrow[d, "q_C"'] & P_{S^C} \arrow[r, two heads, "l_{S^C}"] \arrow[d] & S^C \arrow[d, equal] \\
C \arrow[r, tail, "w^C"] \arrow[d, "d"'] & W^C \arrow[r, two heads] \arrow[d] & S^C \arrow[d, equal] \\
D \arrow[r, tail, "c_d"'] & C_d \arrow[r, two heads, "r"'] & S^C
\end{tikzcd}$$

which induces the following diagram

$$\begin{tikzcd}
\Omega S^C \arrow[r, tail, "k_{S^C}"] \arrow[d, "dq_C"'] & P_{S^C} \arrow[r, two heads, "l_{S^C}"] \arrow[d, "n_{S^C}"] & S^C \arrow[d, equal] \\
D \arrow[r, tail, "c_d"'] & C_d \arrow[r, two heads, "r"'] & S^C.
\end{tikzcd}$$

As P_Q is projective, there exists a morphism $t : P_Q \to P_{S_C}$ such that $l_{S_C} t = r n_Q$.

$$\begin{tikzcd}
 & P_Q \arrow[r, "n_Q"] \arrow[d, dotted, "t"'] & C_d \arrow[d, "r"] \\
\Omega S^C \arrow[r, tail, "k_{S^C}"'] & P_{S^C} \arrow[r, two heads, "l_{S^C}"'] & S^C
\end{tikzcd}$$

Now it follows that $l_{S^C} t k_Q = r n_Q k_Q = r c_d c s_B q_B = 0$, thus there exists a morphism $x : \Omega Q \to \Omega S_C$ such that $k_{S^C} x = t k_Q$.

$$\begin{tikzcd}
\Omega Q \arrow[r, "k_Q"] \arrow[d, dotted, "x"'] & P_Q \arrow[d, "t"] & \\
\Omega S^C \arrow[r, tail, "k_{S_C}"'] & P_{S^C} \arrow[r, two heads, "l_{S^C}"'] & S^C.
\end{tikzcd}$$

As $r n_{S^C} t = l_{S^C} t = r n_Q$, there exists a morphism $y : P_Q \to D$ such that $n_{S^C} t - n_Q = c_d y$. Therefore

$$c_d d q_C x = n_{S^C} k_{S^C} x = n_{S^C} t k_Q = (c_d y + n_Q) k_Q = c_d (y k_Q + c s_B q_B).$$

Then $d q_C x = y k_Q + c s_B q_B$, since c_d is monomorphic. Hence there exists a commutative diagram in $\underline{\mathcal{B}}$

$$\begin{tikzcd}
\Omega Q \arrow[r, "\underline{s_B q_B}"] \arrow[d, "\underline{q_C x}"'] & B \arrow[d, "\underline{c}"] \\
C \arrow[r, "\underline{d}"'] & D.
\end{tikzcd}$$

By Proposition 2.1, we get the following short exact sequences from (6) and (6):

$$\Omega Q \overset{\binom{q_B}{-k_Q}}{\rightarrowtail} \Omega S_B \oplus P_Q \twoheadrightarrow P_{S_B} \oplus S_0, \qquad \Omega S^B \overset{\binom{s_B}{-p}}{\rightarrowtail} B \oplus P_{S^B} \twoheadrightarrow W^B.$$

Then by Proposition 2.2, we get the following commutative diagram of short exact sequences

$$\begin{tikzcd}
\Omega Q \arrow[r, tail, "\binom{q_B}{-k_Q}"] \arrow[d, equal] & \Omega S^B\oplus P_Q \arrow[r, two heads] \arrow[d, tail, "\left(\begin{smallmatrix} s_B & 0\\ -p & 0\\ 0 & 1\end{smallmatrix}\right)"] & P_{S^B}\oplus S_0 \arrow[d, tail] \\
\Omega Q \arrow[r, tail, "\underline{\eta}"'] & B\oplus P_{S^B}\oplus P_Q \arrow[r, two heads] \arrow[d, two heads] & M \arrow[d, two heads] \\
 & W^B \arrow[r, equal] & W^B
\end{tikzcd}$$

where $\underline{\eta}=\underline{s_Bq_B}$. From the third column we get that $M\in\mathcal{U}$. By Lemma 5.5, we obtain that α is epimorphic. □

7. The case where $\underline{\mathcal{H}}$ becomes almost abelian

Definition 7.1. A preabelian category $\mathcal{A}$ is called *left almost abelian* if in any pull-back diagram

$$\begin{tikzcd}
A \arrow[r, "\alpha"] \arrow[d, "\beta"'] & B \arrow[d, "\gamma"] \\
C \arrow[r, "\delta"'] & D
\end{tikzcd}$$

in $\mathcal{A}$, α is a cokernel whenever δ is a cokernel. *Right almost abelian* is defined dually. $\mathcal{A}$ is called *almost abelian* if it is both left and right almost abelian.

In this section we give a sufficient condition when $\underline{\mathcal{H}}$ becomes almost abelian.

We need the following proposition to show our result.

Proposition 7.2. [15, Proposition 2] *Let $A\xrightarrow{f}B\xrightarrow{g}C$ be morphisms in a right (resp. left) semi-abelian category. If f and g are (co-)kernels, then gf is a (co-)kernel. If gf is a (co-)kernel, then f (resp. g) is a (co-)kernel.*

Use this proposition, we can prove the following lemma, which is an analogue of Lemma 5.5

Lemma 7.3. *Let*

$$\begin{tikzcd}
A \arrow[r, "\alpha"] \arrow[d, "\beta"'] & B \arrow[d, "\gamma"] \\
C \arrow[r, "\delta"'] & D
\end{tikzcd}$$

be a pull-back diagram in $\underline{\mathcal{H}}$. If there exists an object $X\in\mathcal{B}^-$ and morphisms $x_B:X\to B$, $x_C:X\to C$ which satisfy that in the following commutative diagram

$$\begin{tikzcd}
X \arrow[r, "\underline{x_B}"] \arrow[d, "\underline{x_C}"'] & B \arrow[d, "\gamma"] \\
C \arrow[r, "\delta"'] & D
\end{tikzcd}$$

$\underline{x_B}$ is a cokernel, then α is a cokernel in $\underline{\mathcal{H}}$.

Proof. Take $x^+:X\to X^+$ as in Definition 3.1. Then by Proposition 3.3, there exist $f_B:X^+\to B$ and $f_C:X^+\to C$ such that $\underline{f_Bx^+}=\underline{x_B}$ and $\underline{f_Cx^+}=\underline{x_C}$. Since $\underline{x_B}$ is a cokernel, by Proposition 7.2, $\underline{f_B}$ is also a cokernel in $\underline{\mathcal{H}}$. As $\gamma\underline{x_B} = \delta\underline{x_C}$, it follows by Proposition 3.3 that $\gamma\underline{f_B} = \delta\underline{f_C}$. By the definition of pull-back, there exists a morphism $\eta : X^+ \to A$ in $\underline{\mathcal{H}}$ which makes the following diagram commute.

$$\begin{tikzcd}
X^+ \arrow[dr, dotted, "\eta"] \arrow[drr, bend left, "\underline{f_B}"] \arrow[ddr, bend right, "\underline{f_C}"'] & & \\
& A \arrow[r, "\alpha"] \arrow[d, "\beta"] & B \arrow[d, "\gamma"] \\
& C \arrow[r, "\delta"'] & D
\end{tikzcd}$$

Since $\underline{f_B}$ is a cokernel, we obtain that α is also epimorphic by Proposition 7.2. □

Theorem 7.4. *Let $(\mathcal{S},\mathcal{T}),(\mathcal{U},\mathcal{V})$ be a twin cotorsion pair on $\mathcal{B}$ satisfying*

$$\mathcal{U} \subseteq \mathcal{T} \text{ or } \mathcal{T} \subseteq \mathcal{U}$$

then $\underline{\mathcal{H}}$ is almost abelian.

Proof. By [15, Proposition 3], a semi-abelian category is left almost abelian if and only if it is right almost abelian. By duality, it is enough to show that $\mathcal{U} \subseteq \mathcal{T}$ implies $\underline{\mathcal{H}}$ is left almost abelian.
Assume we are given a pull-back diagram

$$\begin{tikzcd}
A \arrow[r, "\alpha"] \arrow[d, "\beta"'] & B \arrow[d, "\gamma"] \\
C \arrow[r, "\delta"'] & D
\end{tikzcd}$$

in $\underline{\mathcal{H}}$ where δ is a cokernel. It suffices to show that α becomes a cokernel.
Repeat the same argument as in Theorem 5.6, we get the following diagram

$$\begin{tikzcd}
X \arrow[r, tail, "x_B"] \arrow[d, dotted, "x_C"'] & B \arrow[r, two heads] \arrow[d, dotted, "c"] & S \arrow[d, equal] \\
C \arrow[r, tail, "d"'] & D \arrow[r, two heads] & S.
\end{tikzcd}$$

where $X \in \mathcal{B}^-$, $\underline{d} = \delta$ and $\underline{c} = \gamma$. According to Lemma 7.3, it suffices to show that $\underline{x_B}$ is a cokernel in $\underline{\mathcal{H}}$. By Definition 3.8 and Proposition 2.2, we get the following commutative diagram

$$\begin{tikzcd}
V_B \arrow[r, tail] \arrow[d, equal] & K_{x_B} \arrow[r, two heads, "k_{x_B}"] \arrow[d, tail, "a"] & X \arrow[d, tail, "x_B"] \\
V_B \arrow[r, tail] & W_B \arrow[r, two heads] \arrow[d, two heads] & B \arrow[d, two heads] \\
& S \arrow[r, equal] & S.
\end{tikzcd}$$

It follows that $K_{x_B} \in \mathcal{B}^- = \mathcal{H}$ and $\underline{k_{x_B}x_B} = 0$. By Proposition 5.3, $\underline{x_B}$ is epimorphic in $\underline{\mathcal{H}}$. Now let $r : X \to Q$ be any morphism in $\mathcal{H}$ such that $\underline{rk_{x_B}} = 0$, then rk_{x_B} factors through $\mathcal{W}$. Since $\mathrm{Ext}^1_{\mathcal{B}}(\mathcal{S},\mathcal{T}) = 0$, a is a left $\mathcal{T}$-approximation of K_{x_B}, thus there exists a morphism $b : W_B \to Q$ such that

$ab = rk_B$. By the definition of push-out, we get the following commutative diagram

$$\begin{tikzcd}
K_{x_B} \arrow[r, "k_{x_B}"] \arrow[d, "a"'] & X \arrow[d, "x_B"'] \arrow[rdd, bend left, "r"] & \\
W_B \arrow[r] \arrow[rrd, bend right, "b"'] & B \arrow[rd, dotted] & \\
& & Q.
\end{tikzcd}$$

Since $\underline{x_B}$ is epimorphic in $\underline{\mathcal{H}}$ by Proposition 5.3, the above diagram implies that $\underline{x_B}$ is the cokernel of $\underline{k_{x_B}}$. ☐

By Theorem 6.2, in the case of the above theorem, the heart $\underline{\mathcal{H}}$ also becomes integral. Then by [15, Theorem 2], $\underline{\mathcal{H}}$ is equivalent to a torsionfree class of a hereditary torsion theory in an abelian category induced by $\underline{\mathcal{H}}$. For more details, one can see [15, §4].

8. Existence of enough projectives/injectives

We call an object $P \in \underline{\mathcal{H}}$ (proper-)projective if for any epimorphism (resp. cokernel) $\alpha : X \to Y$ in $\underline{\mathcal{H}}$, there exists an exact sequence

$$\mathrm{Hom}_{\underline{\mathcal{H}}}(P, X) \xrightarrow{\mathrm{Hom}_{\underline{\mathcal{H}}}(P,\alpha)} \mathrm{Hom}_{\underline{\mathcal{H}}}(P, Y) \to 0.$$

An (proper-)injective object is defined dually.
$\underline{\mathcal{H}}$ is said to have enough projectives if for any object $X \in \underline{\mathcal{H}}$, there is a cokernel $\delta : P \to X$ such that P is proper-projective. Having enough injectives is defined dually.

In this section we give sufficient conditions that the heart $\underline{\mathcal{H}}$ of a twin cotorsion pair has enough projectives and has enough injectives.

Lemma 8.1. *If a twin cotorsion pair $(\mathcal{S}, \mathcal{T}), (\mathcal{U}, \mathcal{V})$ satisfies $\mathcal{U} \subseteq \mathcal{T}$, then we have $\Omega\mathcal{S} \subseteq \mathcal{H}$.*

Proof. We first have $\mathcal{P} \subseteq \mathcal{U} = \mathcal{W}$, then by definition $\Omega\mathcal{S} \subseteq \mathcal{B}^-$. But we observe that $\mathcal{U} \subseteq \mathcal{T}$ implies $\mathcal{B}^+ = \mathcal{B}$, hence $\Omega\mathcal{S} \subseteq \mathcal{H}$. ☐

Proposition 8.2. *Let $(\mathcal{S}, \mathcal{T}), (\mathcal{U}, \mathcal{V})$ be a twin cotorsion pair satisfying $\mathcal{U} \subseteq \mathcal{T}$, then any object in $\Omega\mathcal{S}$ is projective in $\underline{\mathcal{H}}$.*

Proof. Let B and C be any objects in $\mathcal{H}$ and let $p : \Omega S \to C$ be any morphism.
Let $\underline{g} : B \to C$ be a morphism which is epimorphic in $\underline{\mathcal{H}}$, by Lemma 5.5 we can assume that it admits a short exact sequence

$$\begin{tikzcd}
A \arrow[r, tail, "f"] & B \arrow[r, two heads, "g"] & C.
\end{tikzcd}$$

Since $B \in \mathcal{H}$ admits a short exact sequence $B \rightarrowtail W^B \twoheadrightarrow S^B$, then according to Proposition 2.2, there exists a commutative diagram

$$\begin{tikzcd}
A \arrow[r, tail, "f"] \arrow[d, equal] & B \arrow[r, two heads, "g"] \arrow[d] & C \arrow[d, "q"] \\
A \arrow[r, tail] & W^B \arrow[r, two heads, "r"] \arrow[d] & D \arrow[d] \\
& S^B \arrow[r, equal] & S^B.
\end{tikzcd}$$

By Lemma 2.10, we obtain $D \in \mathcal{B}^- = \mathcal{H}$. Since $\underline{qg} = 0$ and $\underline{g}$ is epimorphic in $\underline{\mathcal{H}}$, we have $\underline{q} = 0$. By definition ΩS admits a short exact sequence

$$\Omega S \overset{a}{\rightarrowtail} P_S \twoheadrightarrow S \ \ (P \in \mathcal{P}, S \in \mathcal{S}).$$

Since $\underline{qp} = 0$, qp factors through $\mathcal{W}$. As $\mathrm{Ext}^1_{\mathcal{B}}(\mathcal{S}, \mathcal{T}) = 0$, a is a left $\mathcal{T}$-approximation of ΩS. Thus there exists a morphism $s : P \to D$ such that $qp = sa$. Since P is projective, there exists a morphism $t : P \to W^B$ such that $s = rt$. Hence by the definition of pull-back, we get the following commutative diagram

$$\begin{tikzcd}
\Omega S \arrow[rrd, bend left, "p"] \arrow[rd, dotted, "h"] \arrow[rdd, bend right, "ta"'] & & \\
& B \arrow[r, "g"'] \arrow[d] & C \arrow[d, "q"] \\
& W^B \arrow[r, "r"'] & D
\end{tikzcd}$$

which implies that ΩS is projective in $\underline{\mathcal{H}}$. □

Proposition 8.3. *Let* $(\mathcal{S}, \mathcal{T}), (\mathcal{U}, \mathcal{V})$ *be a twin cotorsion pair satisfying* $\mathcal{U} \subseteq \mathcal{T}$, *then any object* $\mathcal{B} \in \mathcal{H}$ *admits an epimorphism* $\alpha : \Omega S \to B$ *in* $\underline{\mathcal{H}}$.

Proof. Let B be any object in $\mathcal{H}$, consider commutative diagram (6). By Proposition 2.1, the left square is a push-out. Now it suffices to show $\underline{s_B}$ is epimorphic in $\underline{\mathcal{H}}$.
Let $c : B \to C$ be any morphism in $\mathcal{H}$ such that $\underline{cs_B} = 0$, then cs_B factors through $\mathcal{W}$. Since p is a left $\mathcal{T}$-approximation of ΩS, there exits a morphism $d : P_{S^B} \to C$ such that $cs_B = dp$. Thus by the definition of push-out we have a commutative diagram

$$\begin{tikzcd}
\Omega S^B \arrow[r, "p"] \arrow[d, "s_B"'] & P_{S^B} \arrow[d] \arrow[rdd, bend left, "d"] & \\
B \arrow[r] \arrow[rrd, bend right, "c"'] & W^B \arrow[rd, dotted] & \\
& & C
\end{tikzcd}$$

which implies $\underline{c} = 0$. Hence $\underline{s_B}$ is epimorphic in $\underline{\mathcal{H}}$. □

Moreover, we have

Proposition 8.4. *Let* $(\mathcal{S}, \mathcal{T}), (\mathcal{U}, \mathcal{V})$ *be a twin cotorsion pair satisfying* $\mathcal{U} \subseteq \mathcal{T}$, *then an object* B *is projective in* $\underline{\mathcal{H}}$ *implies that* $B \in \underline{\Omega \mathcal{S}}$.

Proof. Suppose B is projective in $\underline{\mathcal{H}}$, consider the commutative diagram (6). By Proposition 8.3, $\underline{s_B}$ is epimorphic in $\underline{\mathcal{H}}$, thus B is a direct summand of ΩS^B in $\underline{\mathcal{H}}$. Hence by Lemma 2.9 B lies in $\underline{\Omega \mathcal{S}}$. □

From the following proposition we can get that in the case $\mathcal{U} \subseteq \mathcal{T}$ when the projectives in $\underline{\mathcal{H}}$ is enough.

Proposition 8.5. *Let* $(\mathcal{S}, \mathcal{T}), (\mathcal{U}, \mathcal{V})$ *be a twin cotorsion pair satisfying* $\mathcal{U} \subseteq \mathcal{T}$, *then* $\underline{\mathcal{H}}$ *has enough projectives if and only if any indecomposable object* $B \in \mathcal{H} - \mathcal{U}$ *admits a short exact sequence*

$$B \rightarrowtail S^1 \twoheadrightarrow S^2$$

where $S^1, S^2 \in \mathcal{S}$.

Proof. We prove the "if" part first.
Since an object $B \in \mathcal{H}$ isomorphic to an object $B' \in \mathcal{H}$ in $\underline{\mathcal{H}}$ such that B' does not have any direct

summand in $\mathcal{U}$, we can only consider the obejct $B \in \mathcal{H}$ not having any direct summand in $\mathcal{U}$. Thus by assumption, B admits a short exact sequence

$$B \rightarrowtail S^1 \twoheadrightarrow S^2$$

where $S^1, S^2 \in \mathcal{S}$. As S^2 admits a short exact sequence

$$\begin{tikzcd}\Omega S^2 \arrow[r, tail, "b"] & P_{S^2} \arrow[r, two heads] & S^2.\end{tikzcd}$$

We have the following commutative diagram

$$\begin{tikzcd}
\Omega S^2 \arrow[r, tail, "b"] \arrow[d, "a"'] & P_{S^2} \arrow[r, two heads] \arrow[d] & S^2 \arrow[d, equal] \\
B \arrow[r, tail] & S^1 \arrow[r, two heads] & S^2
\end{tikzcd}$$

Then we get a short exact sequence

$$\begin{tikzcd}\Omega S^2 \arrow[r, tail, "\binom{a}{-b}"] & B \oplus P_{S^2} \arrow[r, two heads] & S^1\end{tikzcd}$$

by Proposition 2.1. Since $B \oplus P_{S^2}$ admits a short exact sequence

$$V \rightarrowtail U \twoheadrightarrow B \oplus P_{S^2}$$

where $V \in \mathcal{V}$ and $U \in \mathcal{U} = \mathcal{W}$, we obtain the following commutative diagram by Proposition 2.2

$$\begin{tikzcd}
V \arrow[r, tail] \arrow[d, equal] & Q \arrow[r, two heads, "c"] \arrow[d, tail, "d"] & \Omega S^2 \arrow[d, tail, "\binom{a}{-b}"] \\
V \arrow[r, tail] & U \arrow[r, two heads] \arrow[d, two heads] & B \oplus P_{S^2} \arrow[d, two heads] \\
& S^1 \arrow[r, equal] & S^1.
\end{tikzcd}$$

Thus $Q \in \mathcal{B}^- = \mathcal{H}$ by Lemma 2.10 and $\underline{ca} = 0$. We claim that $\underline{a}$ is the cokernel of $\underline{c}$ in $\underline{\mathcal{H}}$.
If $r : \Omega S^2 \to M$ is a morphism in $\mathcal{H}$ such that rc factors through $\mathcal{W}$, then there exists $e : U \to M$ such that $cr = ed$, since d is a left $\mathcal{T}$-approximation of Q. Hence by definition of push-out, we get the following commutative diagram

$$\begin{tikzcd}
Q \arrow[r, "c"] \arrow[d] & \Omega S^2 \arrow[d, "\binom{a}{-b}"'] \arrow[ddr, bend left, "r"] & \\
U \arrow[r] \arrow[drr, bend right, "e"'] & B \oplus P_{S^2} \arrow[dr, dotted] & \\
& & M.
\end{tikzcd}$$

which implies that $\underline{r}$ factors through $\underline{a}$. Since $\underline{a}$ is epimorphic in $\underline{\mathcal{H}}$ by Proposition 8.3, we get that $\underline{a}$ is the cokernel of $\underline{c}$.
Now we assume that $\underline{\mathcal{H}}$ has enough projectives.
By Proposition 8.4, all the projective objects in $\underline{\mathcal{H}}$ lie in $\underline{\Omega\mathcal{S}}$. Let B be any indecomposable object in $\mathcal{H} - \mathcal{U}$ and $\beta : \Omega S \to B$ be a cokernel in $\underline{\mathcal{H}}$. Then by Lemma 5.2, we get a short exact sequence

$$\begin{tikzcd}\Omega S \arrow[r, tail, "f"] & B' \arrow[r, two heads] & S'\end{tikzcd}$$

where $B' \in \mathcal{H}$ and $B' \simeq B$ in $\underline{\mathcal{H}}$ and $S' \in \mathcal{S}$. Since ΩS admits a short exact sequence

$$\begin{tikzcd}\Omega S \arrow[r, tail, "p"] & P_S \arrow[r, two heads] & S\end{tikzcd}$$

we take a push-out of f and p, then we get the following commutative diagram

$$\begin{tikzcd}
\Omega S \arrow[r, tail, "f"] \arrow[d, tail, "p"'] & B' \arrow[r, two heads] \arrow[d, tail] & S' \arrow[d, equal] \\
P_S \arrow[r, tail] \arrow[d] & Q' \arrow[r, two heads] \arrow[d] & S' \\
S \arrow[r, equal] & S. &
\end{tikzcd}$$

From the second row we get $Q' \in \mathcal{S}$. Since B is indecomposable, it is a direct summand of B'. Hence by Lemma 2.9, B admits a short exact sequence

$$B \rightarrowtail Q' \twoheadrightarrow S''$$

where $S'' \in \mathcal{S}$. □

By duality, we have

Proposition 8.6. *Let* $(\mathcal{S},\mathcal{T}),(\mathcal{U},\mathcal{V})$ *be a twin cotorsion pair satisfying* $\mathcal{T} \subseteq \mathcal{U}$*, then any monomorphism in* $\underline{\mathcal{H}}$ *is* $\underline{\Omega^{-}\mathcal{V}}$*-monic, and any object in* $\underline{\mathcal{H}}$ *is injective if and only if it lies in* $\underline{\Omega^{-}\mathcal{V}}$*.*

Proposition 8.7. *Let* $(\mathcal{S},\mathcal{T}),(\mathcal{U},\mathcal{V})$ *be a twin cotorsion pair satisfying* $\mathcal{T} \subseteq \mathcal{U}$*, then any object* $\mathcal{B} \in \mathcal{H}$ *admits a monomorphism* $\beta : B \to \Omega^{-}V$ *in* $\underline{\mathcal{H}}$ *where* $\Omega^{-}V \in \underline{\Omega^{-}\mathcal{V}}$*.*

Proposition 8.8. *Let* $(\mathcal{S},\mathcal{T}),(\mathcal{U},\mathcal{V})$ *be a twin cotorsion pair satisfying* $\mathcal{T} \subseteq \mathcal{U}$*, then the heart has enough injectives if and only if any object* $B \in \mathcal{H} - \mathcal{T}$ *admits a short exact sequence*

$$V_2 \rightarrowtail V_1 \twoheadrightarrow B$$

where $V_1, V_2 \in \mathcal{V}$*.*

9. Localisation on the heart of a special twin cotorsion pair

Let $(\mathcal{S},\mathcal{T}),(\mathcal{U},\mathcal{V})$ be a twin cotorsion pair on $\mathcal{B}$ such that $\mathcal{T} = \mathcal{U}$, in this case we get $\mathcal{B}^{+} = \mathcal{B}^{-} = \mathcal{B}$ and $\mathcal{W} = \mathcal{T}$, hence $\underline{\mathcal{H}} = \mathcal{B}/\mathcal{T}$. According to Theorem 6.2, $\mathcal{B}/\mathcal{T}$ is integral. Moreover, By Proposition 8.2 (resp. Proposition 8.6), we obtain that any object in $\Omega\mathcal{S}$ (resp. $\Omega^{-}\mathcal{V}$) is projective (resp. injective) in $\mathcal{B}/\mathcal{T}$.

Let R be the class of regular morphisms in $\mathcal{B}/\mathcal{T}$, then by Theorem [15, p173], the localisation $(\mathcal{B}/\mathcal{T})_R$ (if it exists) is abelian.

Till the end of this section we assume that $\mathcal{B}$ is skeletally small and k-linear over a field k and has a twin cotorsion pair $(\mathcal{S},\mathcal{T}),(\mathcal{T},\mathcal{V})$. We denote that by Proposition 2.12 it is equivalent to assume that $\mathcal{B}$ has a cotorsion pair $(\mathcal{S},\mathcal{T})$ such that $\mathcal{S} \subseteq \mathcal{T}$ and $\mathcal{T}$ is contravariantly finite.

Let $\mathcal{D}$ be a category and R' is a class of morphisms on $\mathcal{D}$. If R' admits both a calculus of right fractions and a calculus of left fractions (for details, see [4, §4]), then the Gabriel-Zisman localisation $\mathcal{D}_{R'}$ at R' (if it exists) has a very nice description. The objects in $\mathcal{D}_{R'}$ are the same as the objects in $\mathcal{D}$. The morphism from X to Y are of the form

$$X \xleftarrow{r} A \xrightarrow{f} Y$$

denoted by $[r, f]$ where r lies in R'.

The localisation functor from $\mathcal{D}$ to $\mathcal{D}_{R'}$ takes a morphism f to $[id, f]$. We denote this image by $[f]$. For $r \in R'$, $[r, id]$ is the inverse of $[r]$. We denote it x_r. Thus, every morphism has the form $[r, f] = [f]x_r$.

By [4, Corollary 4.2], R admits both a calculus of right fractions and a calculus of left fractions.

For a subcategory $\mathcal{C} \subseteq \mathcal{B}$, we denote by $[\mathcal{C}]$ the subcategory of $(\mathcal{B}/\mathcal{T})_R$ which has the same objects as $\mathcal{C}$.

Lemma 9.1. *We have* $\Omega\mathcal{S}/\mathcal{P} = \underline{\Omega\mathcal{S}} \simeq [\Omega\mathcal{S}]$*.*

Proof. We first show that a morphism $f : \Omega S \to B$ factors through $\mathcal{P}$ if and only if it factors through $\mathcal{T}$. Since $\mathcal{P} \subseteq \mathcal{U} = \mathcal{T}$, we only need to show f factors through $\mathcal{T}$ implies it factors through $\mathcal{P}$. Suppose f factors through $\mathcal{T}$. By definition ΩS admits the following short exact sequence

$$\begin{tikzcd}\Omega S \arrow[r, tail, "q"] & P_S \arrow[r, two heads] & S\end{tikzcd}$$

where $P_S \in \mathcal{P}$, $S \in \mathcal{S}$ and B admits the following short exact sequence

$$\begin{tikzcd}V_B \arrow[r, tail] & W_B \arrow[r, two heads, "w_B"] & B.\end{tikzcd}$$

As w_B is a right $\mathcal{U}$-approximation of B, there exists a morphism $a : \Omega S \to W_B$ such that $f = w_B a$. Since q is a left $\mathcal{T}$-approximation of ΩS, there exists a morphism $b : P \to W_B$ such that $bq = a$, hence $f = w_B bq$. Thus by definition we have $\Omega\mathcal{S}/\mathcal{P} = \underline{\Omega\mathcal{S}}$.
Let $L : \underline{\Omega\mathcal{S}} \to [\Omega\mathcal{S}]$ be the location of the localisation functor from $\mathcal{B}/\mathcal{T}$ to $(\mathcal{B}/\mathcal{T})_R$. We claim that it is an equivalence. Obviously it is dense, it is faithful by [4, Lemma 4.4] and full by [4, Lemma 5.4]. $\square$

By this lemma, for convenience, for any morphism f from an object in $\Omega\mathcal{S}$ to $\mathcal{B}$, we also denote its image in $\mathcal{B}/\mathcal{P}$ by $\underline{f}$.

Denote by $\operatorname{Mod}\mathcal{C}$ the category of contravariant additive functors from a category $\mathcal{C}$ to $\operatorname{mod} k$ for any category $\mathcal{C}$. Let $\operatorname{mod}\mathcal{C}$ be the full subcategory of $\operatorname{Mod}\mathcal{C}$ consisting of objects A admitting an exact sequence:

$$\operatorname{Hom}_{\mathcal{C}}(-, C_1) \xrightarrow{\beta} \operatorname{Hom}_{\mathcal{C}}(-, C_0) \xrightarrow{\alpha} A \to 0$$

where $C_0, C_1 \in \mathcal{C}$.

Since $\underline{\Omega\mathcal{S}} \simeq [\Omega\mathcal{S}]$, We have $\operatorname{mod}(\Omega\mathcal{S}/\mathcal{P}) \simeq \operatorname{mod}[\Omega\mathcal{S}]$.

We give the following proposition which is an analogue of [4, Lemma 5.5] (for more details, see [4, §5]).

Proposition 9.2. *If $(\mathcal{S}, \mathcal{T}), (\mathcal{T}, \mathcal{V})$ is a twin cotorsion pair on $\mathcal{B}$ which is skeletally small, and let R denote the class of morphisms which are both monomorphic and epimorphic in $\mathcal{B}/\mathcal{T}$, then*

1. *The projectives in $(\mathcal{B}/\mathcal{T})_R$ are exactly the objects in $\Omega\mathcal{S}$.*
2. *The category $(\mathcal{B}/\mathcal{T})_R$ has enough projectives.*

For convenience, for any objects $X, Y \in \mathcal{B}$, we denote $\operatorname{Hom}_{[\mathcal{B}]}(X, Y)$ by $[X, Y]$. For any morphism $f : X \to Y$, we denote $\operatorname{Hom}_{[\mathcal{B}]}(-, [\underline{f}])$ by $- \circ [\underline{f}]$ and $\operatorname{Hom}_{[\mathcal{B}]}([\underline{f}], -)$ by $[\underline{f}] \circ -$.

Now we can prove the following theorem.

Theorem 9.3. *Let $\mathcal{B}$ be a skeletally small, Krull-Schimdt, k-linear exact category with enough projecitves and injectives, containing a twin cotorsion pair*

$$(\mathcal{S}, \mathcal{T}), (\mathcal{T}, \mathcal{V}).$$

Let R denote the class of morphisms which are both monomorphic and epimorphic in $\mathcal{B}/\mathcal{T}$ and $(\mathcal{B}/\mathcal{T})_R$ denote the localisation of $\mathcal{B}/\mathcal{T}$ at R, then

$$(\mathcal{B}/\mathcal{T})_R \simeq \operatorname{mod}(\Omega\mathcal{S}/\mathcal{P}).$$

Proof. It suffices to show $(\mathcal{B}/\mathcal{T})_R \simeq \operatorname{mod}[\Omega\mathcal{S}]$.
From any object $B \in (\mathcal{B}/\mathcal{T})_R$, there is a projective presentation of B:

$$\Omega S_1 \xrightarrow{[\underline{d_1}]} \Omega S_0 \xrightarrow{[\underline{d_0}]} B \to 0$$

Let ΩS be any object in $[\Omega\mathcal{S}]$, we get the following exact sequence:

$$[\Omega S, \Omega S_1] \xrightarrow{\Omega S \circ [\underline{d_1}]} [\Omega S, \Omega S_0] \xrightarrow{\Omega S \circ [\underline{d_0}]} [\Omega S, B] \to 0$$

which induces a exact sequence in $\operatorname{mod}[\Omega\mathcal{S}]$:

$$[-, \Omega S_1] \xrightarrow{- \circ [\underline{d_1}]} [-, \Omega S_0] \xrightarrow{- \circ [\underline{d_0}]} [-, B] \to 0$$

Now we can define a functor $\Phi : (\mathcal{B}/\mathcal{T})_R \to \operatorname{mod}[\Omega\mathcal{S}]$ as follows:

$$B \mapsto [-, B],$$
$$[\underline{f}] \mapsto - \circ [\underline{f}].$$

• Let us prove that Φ is faithful.
For any morphism $[\underline{f}] : B \to B'$ we have the following commutative diagram

$$\begin{tikzcd}
\Omega S_1 \arrow[r, "{[\underline{d_1}]}"] \arrow[d, dotted, "{[\underline{f_1}]}"'] & \Omega S_0 \arrow[r, "{[\underline{d_0}]}"] \arrow[d, dotted, "{[\underline{f_0}]}"] & B \arrow[r] \arrow[d, "{[\underline{f}]}"] & 0 \\
\Omega S_1' \arrow[r, "{[\underline{d_1'}]}"'] & \Omega S_0 \arrow[r, "{[\underline{d_0'}]}"'] & B \arrow[r] & 0
\end{tikzcd}$$

in $(\mathcal{B}/\mathcal{T})_R$ which induces a commutative diagram in $\operatorname{mod}[\Omega\mathcal{S}]$

$$\begin{tikzcd}
{[-, \Omega S_1]} \arrow[r, "{-\circ[\underline{d_1}]}"] \arrow[d, dotted, "{-\circ[\underline{f_1}]}"'] & {[-, \Omega S_0]} \arrow[r, "{-\circ[\underline{d_0}]}"] \arrow[d, dotted, "{-\circ[\underline{f_0}]}"] & {[-, B]} \arrow[r] \arrow[d, "{-\circ[\underline{f}]}"] & 0 \\
{[-, \Omega S_1']} \arrow[r, "{-\circ[\underline{d_1'}]}"'] & {[-, \Omega S_0']} \arrow[r, "{-\circ[\underline{d_0'}]}"'] & {[-, B']} \arrow[r] & 0.
\end{tikzcd}$$

Hence if $- \circ [\underline{f}] = 0$, we obtain $- \circ [\underline{d_0' f_0}] = 0$, which implies $[\underline{d_0' f_0}] = 0$. Thus $[\underline{f}] = 0$.
• Let us prove that Φ is full.
For any morphism $\alpha : [-, B] \to [-, B']$, we have the following commutative diagram

$$\begin{tikzcd}
{[-, \Omega S_1]} \arrow[r, "{-\circ[\underline{d_1}]}"] \arrow[d, "\alpha_1"'] & {[-, \Omega S_0]} \arrow[r, "{-\circ[\underline{d_0}]}"] \arrow[d, "\alpha_0"'] & {[-, B]} \arrow[r] \arrow[d, "\alpha"] & 0 \\
{[-, \Omega S_1']} \arrow[r, "{-\circ[\underline{d_1'}]}"'] & {[-, \Omega S_0']} \arrow[r, "{-\circ[\underline{d_0'}]}"'] & {[-, B']} \arrow[r] & 0.
\end{tikzcd}$$

in $\operatorname{mod}[\Omega\mathcal{S}]$. By Yoneda's Lemma, there exists $[\underline{f_i}] : \Omega S_i \to \Omega S_i'$ such that $\alpha_i = - \circ [\underline{f_i}]$. Hence there is a commutative diagram

$$\begin{tikzcd}
{[-, \Omega S_1]} \arrow[r, "{-\circ[\underline{d_1}]}"] \arrow[d, "{-\circ[\underline{f_1}]}"'] & {[-, \Omega S_0]} \arrow[r, "{-\circ[\underline{d_0}]}"] \arrow[d, "{-\circ[\underline{f_0}]}"] & {[-, B]} \arrow[r] \arrow[d, dotted, "{-\circ[\underline{f}]}"] & 0 \\
{[-, \Omega S_1']} \arrow[r, "{-\circ[\underline{d_1'}]}"'] & {[-, \Omega S_0']} \arrow[r, "{-\circ[\underline{d_0'}]}"'] & {[-, B']} \arrow[r] & 0.
\end{tikzcd}$$

in $(\mathcal{B}/\mathcal{T})_R$, thus $\alpha = - \circ [\underline{f}]$.
• Let us prove that Φ is dense:
We first show that $\operatorname{mod}[\Omega\mathcal{S}]$ is abelian. It is enough to show that $[\Omega\mathcal{S}]$ has pseudokernels. Let $\alpha : \Omega S_1 \to \Omega S_0$ be a morphism in $[\Omega\mathcal{S}]$, then since $(\mathcal{B}/\mathcal{T})_R$ is abelian, there exists a kernel $\beta : K \to \Omega S_1$ in $(\mathcal{B}/\mathcal{T})_R$. By Proposition 9.2, there exists a epimorphism $\gamma : \Omega S \to K$. We observe that $\beta\gamma$ is a pseudokernel of α. Let $F \in \operatorname{mod}[\Omega\mathcal{S}]$ which admits an exact sequence

$$[-, \Omega S_1] \xrightarrow{-\circ\gamma} [-, \Omega S_0] \to F \to 0$$

where $\gamma \in [\Omega S_1, \Omega S_0]$. Let $B = \operatorname{Coker} \gamma$ then we get an exact sequence

$$\Omega S_1 \xrightarrow{\gamma} \Omega S_0 \to B \to 0$$

in $(\mathcal{B}/\mathcal{T})_R$. Hence $F \simeq [-, B]$. □

10. EXAMPLES

In this section we give several examples of twin cotorsion pair, and we also give some view of the relation between the heart of a cotorsion pair and the hearts of its two components.
First we introduce some notations. Let $\mathcal{C}$ be a subcategory of $\mathcal{B}$, we set

(a) $\mathcal{C}^{\perp_n}=\{X\in\mathcal{B}\mid \operatorname{Ext}^i_{\mathcal{B}}(\mathcal{C},X)=0,\ 0<i\leq n\}$.
(b) ${}^{\perp_n}\mathcal{C}=\{X\in\mathcal{B}\mid \operatorname{Ext}^i_{\mathcal{B}}(X,\mathcal{C})=0,\ 0<i\leq n\}$.
(c) $\mathcal{C}^{\perp}=\{X\in\mathcal{B}\mid \operatorname{Ext}^i_{\mathcal{B}}(\mathcal{C},X)=0,\ \forall i>0\}$.
(d) ${}^{\perp}\mathcal{C}=\{X\in\mathcal{B}\mid \operatorname{Ext}^i_{\mathcal{B}}(X,\mathcal{C})=0,\ \forall i>0\}$.

Definition 10.1. A cotorsion pair $(\mathcal{U},\mathcal{V})$ is called a hereditary cotorsion pair if $\mathcal{U}={}^{\perp}\mathcal{V}$ and $\mathcal{V}=\mathcal{U}^{\perp}$.

Example 10.2. We introduce two trivial hereditary cotorsion pairs:

$$(\mathcal{P},\mathcal{B})\text{ and }(\mathcal{B},\mathcal{I}).$$

We observe that in these two cases the hearts are 0. These two cotorsion pairs also form a twin cotorsion pair

$$(\mathcal{P},\mathcal{B}),(\mathcal{B},\mathcal{I}).$$

We observe that its heart is also 0.

Example 10.3. Let Λ be an artin algebra and T be a cotilting module of finite injective dimension, denote

$$\mathcal{X}:={}^{\perp}T\text{ and }\mathcal{Y}:=({}^{\perp}T)^{\perp}.$$

By [1, Theorem 5.4, Corollary 5.10, Proposition 3.3.], $(\mathcal{X},\mathcal{Y})$ is a hereditary cotorsion pair. By [1, Proposition 3.3, (c, iii)], we get

$$\mathcal{W}\subseteq(\operatorname{mod}\Lambda)^{+}\subseteq\mathcal{Y}.$$

Dually, by [1, Proposition 3.3, (d, iii)], we get

$$\mathcal{W}\subseteq(\operatorname{mod}\Lambda)^{-}\subseteq\mathcal{X}.$$

Then $\mathcal{H}=(\operatorname{mod}\Lambda)^{+}\cap(\operatorname{mod}\Lambda)^{-}\subseteq\mathcal{X}\cap\mathcal{Y}=\mathcal{W}$, hence $\underline{\mathcal{H}}=0$.
By [1, Proposition 1.8], $({}^{\perp_1}T,({}^{\perp_1}T)^{\perp_1})$ is a cotorsion pair. According to [1, §2], ${}^{\perp_1}T,({}^{\perp_1}T)^{\perp_1}$ is also a a cotorsion pair. Hence by definition

$$({}^{\perp}T,({}^{\perp}T)^{\perp}),({}^{\perp_1}T,({}^{\perp_1}T)^{\perp_1})$$

form a twin cotorsion pair. We can also observe that its heart is trivial.

In fact, we have

Proposition 10.4. *If one cotorsion pair in a twin cotorsion pair* $(\mathcal{S},\mathcal{T}),(\mathcal{U},\mathcal{V})$ *is hereditary, then this twin cotorsion pair has a trivial heart, i.e. its heart is zero.*

Proof. We prove that if $(\mathcal{S},\mathcal{T})$ is hereditary, then $\mathcal{W}=\mathcal{B}^{+}\cap\mathcal{B}^{-}=\mathcal{T}\cap\mathcal{S}$, another part is by dual.
For any object $B\in\mathcal{B}^{-}$, there is a short exact category

$$B\rightarrowtail W^B\twoheadrightarrow S^B.$$

Since we have the following exact sequence

$$0=\operatorname{Ext}^1_{\mathcal{B}}(W^B,\mathcal{T})\to\operatorname{Ext}^1_{\mathcal{B}}(B,\mathcal{T})\to\operatorname{Ext}^2_{\mathcal{B}}(S^B,\mathcal{T})=0$$

which implies $B\in\mathcal{S}$. Hence $\mathcal{B}^{-}=\mathcal{S}$. Dually, $\mathcal{B}^{+}=\mathcal{T}$. Hence $\mathcal{W}\subseteq\mathcal{B}^{+}\cap\mathcal{B}^{-}=\mathcal{T}\cap\mathcal{S}\subseteq\mathcal{W}$, this implies $\underline{\mathcal{H}}=0$. □

Recall that $\mathcal{M}$ is *n-cluster tilting* if it satisfies the following conditions

(a) $\mathcal{M}$ is contravariantly finite and covariantly finite in $\mathcal{B}$,
(b) $\mathcal{M}^{\perp_n}=\mathcal{M}$.
(c) ${}^{\perp_n}\mathcal{M}=\mathcal{M}$.

A 2-cluster tilting subcategory is usually called *cluster tilting* subcategory.
Let $\mathcal{M}$ be a cluster tilting subcategory of $\mathcal{B}$. Remark that $\mathcal{P} \subseteq \mathcal{M}$ and $\mathcal{I} \subseteq \mathcal{M}$. For each object $B \in \mathcal{B}$, we have two short exact sequences

$$\begin{tikzcd} B \arrow[r, tail, "f"] & M \arrow[r, two heads] & N, \\ N' \arrow[r, tail] & M' \arrow[r, two heads, "g"] & B \end{tikzcd}$$

that f (resp. g) is a left (resp. right) $\mathcal{M}$-approximation of B. We observe $N \in {}^{\perp_1}\mathcal{M} = \mathcal{M}$ (resp. $N' \in \mathcal{M}^{\perp_1} = \mathcal{M}$), therefore $(\mathcal{M}, \mathcal{M})$ is a cotorsion pair. In this case, $\mathcal{W} = \mathcal{M}$ and $\mathcal{B}^+ = \mathcal{B}^- = \mathcal{B}$, thus $\underline{\mathcal{H}} = \underline{\mathcal{B}} = \mathcal{B}/\mathcal{M}$, which is abelian also by [6].
Moreover, any object in $\Omega\mathcal{M}$ (resp. $\Omega^-\mathcal{M}$) is projective (resp. injective) in $\mathcal{B}/\mathcal{M}$, and by Proposition 8.5,8.8, $\mathcal{B}/\mathcal{M}$ has enough projectives and enough injectives.

Proposition 10.5. *A subcategory $\mathcal{M}$ in $\mathcal{B}$ is cluster tilting if and only if $(\mathcal{M}, \mathcal{M})$ is a cotorsion pair on $\mathcal{B}$.*

Proof. From the above discussion, we know that $(\mathcal{M}, \mathcal{M})$ is a cotorsion pair if $\mathcal{M}$ is cluster tilting, so it remains to show the "only if" part. But it is just followed by the definition of cotorsion pair and Lemma 2.4. ☐

In the following examples, we denote by "$\circ$" in a quiver the objects belong to a subcategory and by "$\cdot$" the objects do not.

Example 10.6. Let Λ be the path algebra of the following quiver

$$\begin{tikzcd} 1 & 2 \arrow[l] & 3 \arrow[l] & 4 \arrow[l] \arrow[lll, dotted, bend right] \end{tikzcd}$$

then we obtain the AR-quiver $\Gamma(\operatorname{mod}\Lambda)$ of $\operatorname{mod}\Lambda$.



Let $\mathcal{M} = \{X \in \operatorname{mod}\Lambda \mid \operatorname{Ext}^1_{\mathcal{B}}(X, \Lambda) = 0\}$, then by [1, Proposition 1.10, 1.9], $(\mathcal{M}, \mathcal{M}^{\perp_1})$ is a cotorsion pair on $\operatorname{mod}\Lambda$. But

$$\mathcal{M} = \begin{array}{ccccccc} \circ & & \cdot & & \cdot & & \circ \\ & \circ & & \cdot & & \circ & \\ & & \circ & & \circ & & \end{array}$$

which consisting of all the direct sums of indecomposable projectives and indecomposable injectives. We observe that in fact $\mathcal{M} = \mathcal{M}^{\perp_1}$ and hence it is a cluster tilting subcategory. And the quiver of the quotient category $\operatorname{mod}\Lambda/\mathcal{M}$ is



which is equivalent to the AR-quiver of A_2.

Example 10.7. Take the notion of the former example, Let

$$\mathcal{M}' = \begin{array}{ccccccc} \circ & & \cdot & & \cdot & & \cdot \\ & \circ & & \cdot & & \circ & \\ & & \circ & & \circ & & \end{array}$$

then by [1, Proposition 1.10, 1.9], $(\mathcal{M}', \mathcal{M}'^{\perp_1})$ is a cotorsion pair and

$$\mathcal{M}'^{\perp_1} = \begin{matrix} \circ & & \cdot & & \circ & & \circ \\ & \circ & & \cdot & & \circ & \\ & & \circ & & \circ & & \end{matrix}$$

hence it contains Λ. Obviously it is closed under extension and contravariantly finite, then by [1, Proposition 1.10, 1.9], $(\mathcal{M}'^{\perp_1}, (\mathcal{M}'^{\perp_1})^{\perp_1})$ is also a cotorsion pair on $\operatorname{mod}\Lambda$ and

$$(\mathcal{M}'^{\perp_1})^{\perp_1} = \begin{matrix} \circ & & \cdot & & \cdot & & \circ \\ & \cdot & & \cdot & & \circ & \\ & & \circ & & \circ & & \end{matrix}$$

Thus we get a twin cotorsion pair

$$(\mathcal{M}', \mathcal{M}'^{\perp_1}), (\mathcal{M}'^{\perp_1}, (\mathcal{M}'^{\perp_1})^{\perp_1}).$$

Then the quiver of $\operatorname{mod}\Lambda/\mathcal{M}'^{\perp_1}$ is $2 \to \begin{smallmatrix}3\\2\end{smallmatrix}$. The quiver of quotient category $\Omega\mathcal{M}'/\mathcal{P}$ is just 2. Hence we get $(\operatorname{mod}\Lambda/\mathcal{M}'^{\perp_1})_R \simeq \operatorname{mod}(\Omega\mathcal{M}'/\mathcal{P})$.

By Proposition 10.4, the heart of a twin cotorsion pair can become zero when one heart of its component is zero. From the following example we can see that there are also some conditions in which the heart of a twin cotorsion pair is non-trivial when only one heart of its components is zero.

Example 10.8. Take the notion of Exmaple 10.6, let

$$\mathcal{S} = \begin{matrix} \circ & & \cdot & & \circ & & \cdot \\ & \circ & & \cdot & & \cdot & \\ & & \circ & & \circ & & \end{matrix} \quad \mathcal{T} = \begin{matrix} \circ & & \cdot & & \circ & & \circ \\ & \cdot & & \circ & & \circ & \\ & & \circ & & \circ & & \end{matrix}$$

and

$$\mathcal{U} = \begin{matrix} \circ & & \cdot & & \circ & & \circ \\ & \circ & & \cdot & & \circ & \\ & & \circ & & \circ & & \end{matrix} \quad \mathcal{V} = \begin{matrix} \circ & & \cdot & & \cdot & & \circ \\ & \cdot & & \cdot & & \circ & \\ & & \circ & & \circ & & \end{matrix}$$

then $(\mathcal{S}, \mathcal{T})$ and $(\mathcal{U}, \mathcal{V})$ are cotorsion pairs and $(\mathcal{S}, \mathcal{T})(\mathcal{U}, \mathcal{T})$ is a twin cotorsion pair. The heart of $(\mathcal{S}, \mathcal{T})$ is zero and the heart of $(\mathcal{U}, \mathcal{V})$ is $\operatorname{add}(\begin{smallmatrix}3\\2\end{smallmatrix})$. Now we have

$$\mathcal{W} = \mathcal{U} \cap \mathcal{T} = \begin{matrix} \circ & & \cdot & & \circ & & \circ \\ & \cdot & & \cdot & & \circ & \\ & & \circ & & \circ & & \end{matrix}$$

then we get

$$(\operatorname{mod}\Lambda)^+/\mathcal{W} = \operatorname{add}(\begin{smallmatrix}3\\2\end{smallmatrix}) \text{ and } (\operatorname{mod}\Lambda)^-/\mathcal{W} = \operatorname{add}(\begin{smallmatrix}2\\1\end{smallmatrix} \oplus 2 \oplus \begin{smallmatrix}3\\2\end{smallmatrix})$$

Hence $\underline{\mathcal{H}} = \operatorname{add}(\begin{smallmatrix}3\\2\end{smallmatrix})$.
One can check that the twin cotorsion pair $(\mathcal{S}, \mathcal{T}), (\mathcal{U}, \mathcal{V})$ where

$$\mathcal{S} = \begin{matrix} \circ & & \cdot & & \cdot & & \circ \\ & \circ & & \cdot & & \cdot & \\ & & \circ & & \circ & & \end{matrix} \quad \mathcal{T} = \begin{matrix} \circ & & \circ & & \cdot & & \circ \\ & \circ & & \cdot & & \circ & \\ & & \circ & & \circ & & \end{matrix}$$

and

$$\mathcal{U} = \begin{matrix} \circ & & \circ & & \cdot & & \circ \\ & \circ & & \circ & & \cdot & \\ & & \circ & & \circ & & \end{matrix} \quad \mathcal{V} = \begin{matrix} \cdot & & \circ & & \cdot & & \circ \\ & \cdot & & \cdot & & \circ & \\ & & \circ & & \circ & & \end{matrix}$$

gives an example of another condition.

From Example 10.7, we see that there exits two cotorsion pair which have non-trivial hearts form a twin cotorsion pair also having a non-trivial heart. From the following example, we see that even two components of a twin cotorsion pair have non-trivial hearts, the heart of the twin cotorsion pair itself can be zero.

Example 10.9. Let Λ be the k-algebra given by the quiver



and bound by $\alpha\beta = 0$ and $\beta\gamma\alpha = 0$. Then its AR-quiver $\Gamma(\operatorname{mod}\Lambda)$ is given by



Here, the first and the last columns are identified. Let



and



The heart of cotorsion pair $(\mathcal{S},\mathcal{T})$ is add(1) and the heart of cotorsion pair $(\mathcal{U},\mathcal{V})$ is add(3). But when we consider the twin cotorsion pair $(\mathcal{S},\mathcal{T}),(\mathcal{U},\mathcal{V})$, we get $\mathcal{W} = \mathcal{V}$ and

$$(\operatorname{mod}\Lambda)^{-}/\mathcal{W} = \operatorname{add}(\,1 \oplus 2\,) \text{ and } (\operatorname{mod}\Lambda)^{+}/\mathcal{W} = \operatorname{add}(\,3\,)$$

hence its heart is zero.

Acknowledgments

The author would like to thank Laurent Demonet and Osamu Iyama for their helpful advices and corrections.

References

[1] M. Auslander, I. Reiten. Application of contravariantly finite subcategories. Adv. Math. 86, 111–152(1991).
[2] T. Bühler. Exact categories. Expo. Math. 28 (2010), no. 1, 1–69.
[3] A. A. Beilinson, J. Bernstein, P. Deligne. Faisceaux pervers. Analysis and topology on singular spaces, I (Luminy 1981), 5–171, Astérisque, 100, Soc. Math. France, Pairs, 1982.
[4] A. B. Buan, R. J. Marsh. From triangulated categories to module categories via localisation II: calculus of fractions. arXiv: 1102.4597v2.
[5] R. R. Colby, K. R. Fuller. Tilting, cotilting, and serially tilted rings. Comm. in Alg. 17 (1990), 1709–1722.
[6] L. Demonet, Y. Liu. Quotients of exact categories by cluster tilting subcategories as module categories. arXiv: 1208.0639.
[7] Z. Huang, O. Iyama. Auslander-type conditions and cotorsion pairs. J. Algebra. 318 (2007), 93–110.
[8] O. Iyama. Cluster tilting for higher Auslander algebra. Adv. Math. 226 (2011), no. 1, 1–61.
[9] O. Iyama, Y. Yoshino. Mutation in triangulated categoryies and rigid Cohen-Macaulay modules Invent. Math. 172 (2008), no. 1, 117–168.
[10] B. Keller. Chain complexes and stable categories. Manuscripta Math. 67 (1990), no. 4, 379–417.
[11] H. Krause, Ø. Solberg. Application of cotorsion pairs. J. London Math. Soc. (2) 68 (2003) 631–650.

[12] S. Koenig, B. Zhu. From triangulated categories to abelian categories: cluster tilting in a general framework. Math. Z. 258 (2008), no. 1, 143–160.

[13] H. Nakaoka. General heart construction on a triangulated category (I): unifying t-structures and cluster tilting subcategories. Appl. Categ. Structures 19 (2011), no. 6, 879–899.

[14] H. Nakaoka. General heart construction for twin torsion pairs on triangulated categories. arXiv: 1111.1820.

[15] W. Rump. Almost abelian categories. Cahiers Topologie Géom. Différentielle. Catég. 42 (2003), no. 3, 163–225.

[16] L. Salce. Cotorsion theories for abelian groups. Symposia Mathmatica 23 (Cambridge University Press, Cambridge, 1979) 11–32

Graduate School of Mathematics, Nagoya University, 464-8602 Nagoya, Japan
E-mail address: `d11005m@math.nagoya-u.ac.jp`